# antibodies .- online.com





# CD43 Protein (AA 20-395) (rho-1D4 tag)



## **Image**



#### Go to Product page

#### Overview

Quantity:	1 mg
Target:	CD43 (SPN)
Protein Characteristics:	AA 20-395
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD43 protein is labelled with rho-1D4 tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA, Crystallization (Crys)

#### **Product Details**

### Sequence:

DSLQRTTMLP STPHITAPST SEAQNASPSV SVGSGTVDSK ETISPWGQTT IPVSLTPLET
TELSSLETSA GASMSTPVPE PTASQEVSSK TSALLPEPSN VASDPPVTAA NPVTDGPAAN
PVTDGTAAST SISKGTSAPP TTVTTSSNET SGPSVATTVS SKTSGPPVTT ATGSLGPSSE
MHGLPATTAT SSVESSSVAR GTSVSSRKTS TTSTQDPITT RSPSQESSGM LLVPMLIALV
VVLALVALLL LWRQRQKRRT GALTLSGGGK RNGVVDAWAG PARVPDEEAT TTSGAGGNKG
SEVLETEGSG QRPTLTTFFS RRKSRQGSLV LEELKPGSGP NLKGEEEPLV GSEDEAVETP
TSDGPQAKDE AAPQSL

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

#### Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Spn Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

- 1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

 Purity:
 >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.


 Sterility:
 0.22 μm filtered

 Endotoxin Level:
 Protein is endotoxin-free.

 Grade:
 Crystallography grade

## **Target Details**

of the cell and allow multiple glycan chains to be accessible for binding. Is a counter-receptor for SN/Siglec-1. During T-cell activation is actively removed from the T-cell-APC (antigen-presenting cell) contact site thus suggesting a negative regulatory role in adaptive immune response. {ECO:0000269 PubMed:11728336}.  39.2 kDa Including tag.
physicochemical properties of the T-cell surface and in lectin binding. Presents carbohydrate ligands to selectins. Has an extended rodlike structure that could protrude above the glycocalyx of the cell and allow multiple glycan chains to be accessible for binding. Is a counter-receptor for SN/Siglec-1. During T-cell activation is actively removed from the T-cell-APC (antigen-presenting cell) contact site thus suggesting a negative regulatory role in adaptive immune response. {ECO:0000269 PubMed:11728336}.
ligands to selectins. Has an extended rodlike structure that could protrude above the glycocalyx of the cell and allow multiple glycan chains to be accessible for binding. Is a counter-receptor for SN/Siglec-1. During T-cell activation is actively removed from the T-cell-APC (antigen-presenting cell) contact site thus suggesting a negative regulatory role in adaptive immune response. {ECO:0000269 PubMed:11728336}.  39.2 kDa Including tag.
for SN/Siglec-1. During T-cell activation is actively removed from the T-cell-APC (antigen-presenting cell) contact site thus suggesting a negative regulatory role in adaptive immune response. {ECO:0000269 PubMed:11728336}.  39.2 kDa Including tag.
for SN/Siglec-1. During T-cell activation is actively removed from the T-cell-APC (antigen-presenting cell) contact site thus suggesting a negative regulatory role in adaptive immune response. {ECO:0000269 PubMed:11728336}.  39.2 kDa Including tag.
presenting cell) contact site thus suggesting a negative regulatory role in adaptive immune response. {ECO:0000269 PubMed:11728336}.  39.2 kDa Including tag.
response. {ECO:0000269 PubMed:11728336}.  39.2 kDa Including tag.
39.2 kDa Including tag.
D1 F700
P15702
Regulation of Leukocyte Mediated Immunity
In addition to the applications listed above we expect the protein to work for functional studies
as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
though.
Protein has not been tested for activity yet. In cases in which it is highly likely that the
recombinant protein with the default tag will be insoluble our protein lab may suggest a higher
molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible
options with you in detail to assure that you receive your protein of interest.
For Research Use only
Liquid
100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Avoid repeated freeze-thaw cycles.
-80 °C
Store at -80°C.
Unlimited (if stored properly)



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process