antibodies .- online.com





FGFR1 Protein (AA 398-822) (His tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	FGFR1
Protein Characteristics:	AA 398-822
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FGFR1 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:

KMKSGTKKSD FHSQMAVHKL AKSIPLRRQV TVSADSSASM NSGVLLVRPS RLSSSGTPML
AGVSEYELPE DPRWELPRDR LVLGKPLGEG CFGQVVLAEA IGLDKDKPNR VTKVAVKMLK
SDATEKDLSD LISEMEMMKM IGKHKNIINL LGACTQDGPL YVIVEYASKG NLREYLQARR
PPGLEYCYNP SHNPEEQLSS KDLVSCAYQV ARGMEYLASK KCIHRDLAAR NVLVTEDNVM
KIADFGLARD IHHIDYYKKT TNGRLPVKWM APEALFDRIY THQSDVWSFG VLLWEIFTLG
GSPYPGVPVE ELFKLLKEGH RMDKPSNCTN ELYMMMRDCW HAVPSQRPTF KQLVEDLDRI
VALTSNQEYL DLSIPLDQYS PSFPDTRSST CSSGEDSVFS HEPLPEEPCL PRHPTQLANS GLKRR
Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Fgfr1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 μm filtered
Endotoxin Level:	Protein is endotoxin free.
Grade:	Crystallography grade

Target Details

Target:	FGFR1
Alternative Name:	Fgfr1 (FGFR1 Products)

Target Details

Background:	Tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and
	plays an essential role in the regulation of embryonic development, cell proliferation,
	differentiation and migration. Required for normal mesoderm patterning and correct axial
	organization during embryonic development, normal skeletogenesis and normal development
	of the gonadotropin-releasing hormone (GnRH) neuronal system. Phosphorylates PLCG1, FRS2,
	GAB1 and SHB. Ligand binding leads to the activation of several signaling cascades. Activation
	of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol
	1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and
	SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase
	signaling pathway, as well as of the AKT1 signaling pathway. Promotes phosphorylation of
	SHC1, STAT1 and PTPN11/SHP2. In the nucleus, enhances RPS6KA1 and CREB1 activity and
	contributes to the regulation of transcription. FGFR1 signaling is down-regulated by
	IL17RD/SEF, and by FGFR1 ubiquitination, internalization and degradation (By similarity).
	{ECO:0000250 UniProtKB:P11362, ECO:0000269 PubMed:10821861,
	ECO:0000269 PubMed:10896947, ECO:0000269 PubMed:1309590,
	ECO:0000269 PubMed:17086194, ECO:0000269 PubMed:8001822,
	ECO:0000269 PubMed:8001823}.
Molecular Weight:	48.8 kDa Including tag.
UniProt:	P16092
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin
	Signaling Pathway, Sensory Perception of Sound, Stem Cell Maintenance, S100 Proteins
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the
	recombinant protein with the default tag will be insoluble our protein lab may suggest a higher
	molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible
	options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process