antibodies .- online.com





POU4F1 Protein (AA 1-421) (Strep Tag)



Go to Product pag

Overview

Quantity:	1 mg
Target:	POU4F1
Protein Characteristics:	AA 1-421
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This POU4F1 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence:

MMSMNSKQPH FAMHPTLPEH KYPSLHSSSE AIRRACLPTP PLQSNLFASL DETLLARAEA LAAVDIAVSQ GKSHPFKPDA TYHTMNSVPC TSTSTVPLAH HHHHHHHHHQA LEPGDLLDHI SSPSLALMAG AGGAGAAGGG GGAHDGPGGG GGPGGGGGPG GGGPGGGGGG GGPGGGGGP GGGLLGGSAH PHPHMHGLGH LSHPAAAAAM NMPSGLPHPG LVAAAAHHGA AAAAAAAAAG QVAAASAAAA VVGAAGLASI CDSDTDPREL EAFAERFKQR RIKLGVTQAD VGSALANLKI PGVGSLSQST ICRFESLTLS HNNMIALKPI LQAWLEEAEG AQREKMNKPE LFNGGEKKRK RTSIAAPEKR SLEAYFAVQP RPSSEKIAAI AEKLDLKKNV VRVWFCNQRQ KQKRMKFSAT Y Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Product Details

Purity:	≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Endotoxin Levei:	Low Endotoxin less than T EU/mg (< 0.1 ng/mg)
Target Details	
Target:	POU4F1
Alternative Name:	Pou4f1 (POU4F1 Products)
Background:	POU domain, class 4, transcription factor 1 (Brain-specific homeobox/POU domain protein 3A)
	(Brain-3A) (Brn-3A) (Brn-3.0),FUNCTION: Multifunctional transcription factor with different
	regions mediating its different effects (PubMed:10640682, PubMed:8621561,
	PubMed:9694219, PubMed:9722627). Acts by binding (via its C-terminal domain) to sequences
	related to the consensus octamer motif 5'-ATGCAAAT-3' in the regulatory regions of its target
	genes (PubMed:8621561, PubMed:17668438). Regulates the expression of specific genes
	involved in differentiation and survival within a subset of neuronal lineages. It has been shown
	that activation of some of these genes requires its N-terminal domain, maybe through a
	neuronal-specific cofactor (PubMed:12934100). Ativates BCL2 expression and protects
	neuronal cells from apoptosis (via the N-terminal domain) (PubMed:9722627). Induces
	neuronal process outgrowth and the coordinate expression of genes encoding synaptic
	proteins (PubMed:8972215). Exerts its major developmental effects in somatosensory neurons
	and in brainstem nuclei involved in motor control. Stimulates the binding affinity of the nuclear
	estrogene receptor ESR1 to DNA estrogen response element (ERE), and hence modulates
	ESR1-induced transcriptional activity (PubMed:9448000). May positively regulate POU4F2 and
	POU4F3 (PubMed:8876243). Regulates dorsal root ganglion sensory neuron specification and
	axonal projection into the spinal cord (PubMed:22326227). Plays a role in TNFSF11-mediated
	terminal osteoclast differentiation (PubMed:17668438). Negatively regulates its own
	expression interacting directly with a highly conserved autoregulatory domain surrounding the
	transcription initiation site (PubMed:12441296). {ECO:0000269 PubMed:10640682,
	ECO:0000269 PubMed:12441296, ECO:0000269 PubMed:12934100,
	ECO:0000269 PubMed:17668438, ECO:0000269 PubMed:22326227,
	ECO:0000269 PubMed:8621561, ECO:0000269 PubMed:8876243,
	ECO:0000269 PubMed:8972215, ECO:0000269 PubMed:9448000,
	ECO:0000269 PubMed:9694219, ECO:0000269 PubMed:9722627}., FUNCTION: [Isoform 2]:
	Able to act as transcription factor, cannot regulate the expression of the same subset of genes
	than isoform 1 (PubMed:12934100). Does not have antiapoptotic effect on neuronal cells

(PubMed:9722627). {ECO:0000269|PubMed:12934100, ECO:0000269|PubMed:9722627}.

Target Details

Molecular Weight:	42.8 kDa
UniProt:	P17208
Pathways:	Feeding Behaviour

·	
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional
	components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's

needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)