

Datasheet for ABIN3133309

**FGFR2 Protein (AA 22-377) (His tag)**[Go to Product page](#)**1** Image

## Overview

Quantity:	1 mg
Target:	FGFR2
Protein Characteristics:	AA 22-377
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FGFR2 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

## Product Details

Sequence:	<p>RPSFSLVEDT TLEPEEPPTK YQISQPEAYV VAPGESLELQ CMLKDAAVIS WTKDGVHLGP NNRTVLIGEY LQIKGATPRD SGLYACTAAR TVDSETWIFM VNVTDAISSG DDEDDTDSSE DVVSENRSNQ RPYWTNTEK MEKRLHACPA ANTVKFRCPA GGNPTSTMRW LKNGKEFKQE HRIGGYKVRN QHWSLIMESV VPSDKGNYTC LVENEYGSIN HTYHLDVVER SPHRPILQAG LPANASTVVG GDVEFVCKVY SDAQPHIQWI KHVEKNGSKN GPDGLPYLKV LKAAGVNTTD KEIEVLYIRN VTFEDAGEYT CLAGNSIGIS FHSAWLTVLP APVREKEITA SPDYLE</p> <p><b>Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.</b></p>
Characteristics:	<ul style="list-style-type: none"><li>• Made in Germany - from design to production - by highly experienced protein experts.</li><li>• Mouse Fgfr2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.</li><li>• State-of-the-art algorithm used for plasmid design (Gene synthesis).</li></ul>

## Product Details

---

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the ExPASy's protparam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in baculovirus infected SF9 insect cells: <ol style="list-style-type: none"><li>1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li><li>2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li></ol>
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Protein is endotoxin free.
Grade:	Crystallography grade

## Target Details

---

Target:	FGFR2
Alternative Name:	Fgfr2 ( <a href="#">FGFR2 Products</a> )
Background:	Tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and

## Target Details

plays an essential role in the regulation of cell proliferation, differentiation, migration and apoptosis, and in the regulation of embryonic development. Required for normal embryonic patterning, trophoblast function, limb bud development, lung morphogenesis, osteogenesis and skin development. Plays an essential role in the regulation of osteoblast differentiation, proliferation and apoptosis, and is required for normal skeleton development. Promotes cell proliferation in keratinocytes and immature osteoblasts, but promotes apoptosis in differentiated osteoblasts. Phosphorylates PLCG1, FRS2 and PAK4. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. FGFR2 signaling is down-regulated by ubiquitination, internalization and degradation. Mutations that lead to constitutive kinase activation or impair normal FGFR2 maturation, internalization and degradation lead to aberrant signaling. Over-expressed FGFR2 promotes activation of STAT1. {ECO:0000269|PubMed:10851026, ECO:0000269|PubMed:15629145, ECO:0000269|PubMed:8393815, ECO:0000269|PubMed:8663044}.

Molecular Weight: 40.4 kDa Including tag.

UniProt: [P21803](#)

Pathways: [RTK Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Regulation of Muscle Cell Differentiation](#), [Skeletal Muscle Fiber Development](#), [Growth Factor Binding](#)

## Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

## Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

## Images



**Image 1.** „Crystallography Grade“ protein due to multi-step, protein-specific purification process