

# Datasheet for ABIN3133676

# SSTR2 Protein (AA 1-369) (Strep Tag)



$\sim$			
( )\	<b>/</b> e	rVI	iew

Quantity:	250 μg
Target:	SSTR2
Protein Characteristics:	AA 1-369
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SSTR2 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details	
Brand:	AliCE®
Sequence:	MEMSSEQLNG SQVWVSSPFD LNGSLGPSNG SNQTEPYYDM TSNAVLTFIY FVVCVVGLCG
	NTLVIYVILR YAKMKTITNI YILNLAIADE LFMLGLPFLA MQVALVHWPF GKAICRVVMT
	VDGINQFTSI FCLTVMSIDR YLAVVHPIKS AKWRRPRTAK MINVAVWCVS LLVILPIMIY
	AGLRSNQWGR SSCTINWPGE SGAWYTGFII YAFILGFLVP LTIICLCYLF IIIKVKSSGI
	RVGSSKRKKS EKKVTRMVSI VVAVFIFCWL PFYIFNVSSV SVAISPTPAL KGMFDFVVIL
	TYANSCANPI LYAFLSDNFK KSFQNVLCLV KVSGTEDGER SDSKQDKSRL NETTETQRTL
	LNGDLQTSI
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- · State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made
Target Details	
Target:	SSTR2

## **Target Details**

Alternative Name:	Sstr2 (SSTR2 Products)
Background:	Somatostatin receptor type 2 (SS-2-R) (SS2-R) (SS2R) (SRIF-1),FUNCTION: Receptor for
	somatostatin-14 and -28. This receptor is coupled via pertussis toxin sensitive G proteins to
	inhibition of adenylyl cyclase. In addition it stimulates phosphotyrosine phosphatase and PLC
	via pertussis toxin insensitive as well as sensitive G proteins. Inhibits calcium entry by
	suppressing voltage-dependent calcium channels. Acts as the functionally dominant
	somatostatin receptor in pancreatic alpha- and beta-cells where it mediates the inhibitory effect
	of somatostatin-14 on hormone secretion. Inhibits cell growth through enhancement of MAPK
	and MAPK2 phosphorylation and subsequent up-regulation of CDKN1B. Stimulates neuronal
	migration and axon outgrowth and may participate in neuron development and maturation
	during brain development. Mediates negative regulation of insulin receptor signaling through
	PTPN6. Inactivates SSTR3 receptor function following heterodimerization.
	{ECO:0000269 PubMed:8104154, ECO:0000269 PubMed:9507021}.
Molecular Weight:	41.2 kDa
UniProt:	P30875
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	
	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce
Restrictions:	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's

## Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.  Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol <b>Might differ depending on protein.</b>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months