

Datasheet for ABIN3134236 ERCC3 Protein (AA 1-783) (Strep Tag)



Overview

Quantity:	250 μg
Target:	ERCC3
Protein Characteristics:	AA 1-783
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This ERCC3 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA	
Product Details		
Brand:	AliCE®	
Sequence:	MGKRDRVDRD KKKSKKRQYE EEEEDEDDIP GNESQEAVPS AAGKQVDESS TKVDEYGAKD	
	YRQQMPLKGD HTSRPLWVAP DGHIFLEAFS PVYKYAQDFL VAIAEPVCRP THVHEYKLTA	
	YSLYAAVSVG LQTSDITEYL RKLSKTGVPD GIIQFIKLCT VSYGKVKLVL KHNRYFVESS	
	HPDVIQHLLQ DPVIRECRLR NAEGEATELI TETFTSKSAI SKTAAEGSGG PSTSQGVDAQ	
	ATSDIPKDLF DFYEQMDKDE EEEEETQTVS FEVKQEMIEE LQKRCICLEY PLLAEYDFRN	
	DTLNPDINID LKPTAVLRPY QEKSLRKMFG NGRARSGVIV LPCGAGKSLV GVTAACTVRK	
	RCLVLGNSAV SVEQWKAQFK MWSTIDDSQI CRFTSDAKDK PIGCSVAIST YSMLGHTTKR	
	SWEAERVMEW LKTQEWGLMI LDEVHTIPAR MFRRVLTIVQ AHCKLGLTAT LVREDDKIVD	
	LNFLIGPKLY EANWMELQNN GYIAKVQCAE VWCPMSPEFY REYVAIKTKK RILLYTMNPN	
	KFRACQFLIK FHERRNDKII VFADNVFALK EYAIRLNKPY IYGPTSQGER MQILQNFKHN	
	PKINTIFISK VGDTSFDLPE ANVLIQISSH GGSRRQEAQR LGRVLRAKKG MVAEEYNAFF	

YSLVSQDTQE MAYSTKRQRF LVDQGYSFKV ITKLAGMEEE ELAFSTKEEQ QQLLQKVLAA
TDLDAEEEVV AGEFGSRSGQ ASRRCGTMSS LSGADDTVYM EYHSSRSKAS SKHVHPLFKR FRK

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).

Product Details	
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made
Target Details	
Target:	ERCC3
Alternative Name:	Ercc3 (ERCC3 Products)
Background:	General transcription and DNA repair factor IIH helicase subunit XPB (TFIIH subunit XPB) (EC
	3.6.4.12) (Basic transcription factor 2 89 kDa subunit) (BTF2 p89) (DNA excision repair protein
	ERCC-3) (DNA repair protein complementing XP-B cells) (TFIIH 89 kDa subunit) (Xeroderma
	pigmentosum group B-complementing protein),FUNCTION: ATP-dependent 3'-5' DNA helicase
	component of the general transcription and DNA repair factor IIH (TFIIH) core complex, which
	is involved in general and transcription-coupled nucleotide excision repair (NER) of damaged
	DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II. In NER, TFIIH
	acts by opening DNA around the lesion to allow the excision of the damaged oligonucleotide
	and its replacement by a new DNA fragment. The ATPase activity of XPB/ERCC3, but not its
	helicase activity, is required for DNA opening. In transcription, TFIIH has an essential role in
	transcription initiation. When the pre-initiation complex (PIC) has been established, TFIIH is
	required for promoter opening and promoter escape. The ATP-dependent helicase activity of
	XPB/ERCC3 is required for promoter opening and promoter escape. Phosphorylation of the C-
	terminal tail (CTD) of the largest subunit of RNA polymerase II by the kinase module CAK
	controls the initiation of transcription. {ECO:0000250 UniProtKB:P19447}.
Molecular Weight:	89.1 kDa
UniProt:	P49135
Pathways:	DNA Damage Repair
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce

even the most difficult-to-express proteins, including those that require post-translational

Application Details

modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months