

Datasheet for ABIN3134516

CIDEc Protein (AA 1-239) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	CIDEc
Protein Characteristics:	AA 1-239
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CIDEc protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Brand:	ALiCE®
Sequence:	MDYAMKSLSL LYPRSLSRHV AVSTAVVTQQ LVSKPSRETP RARPCRVSTA DRKVRKGIMA HSLEDLLNKV QDILKLKDKP FSLVLEEDGT IVETEEYFQA LAKDTMFMVL LKGQKWKPPS EQRKKRAQLA LSQKPTKKID VARVTFDLYK LNPQDFIGCL NVKATLYDTY SLSYDLHCYK AKRIVKEMLR WTLFSMQATG HMLLGTSSYM QQFLDATEEE QPAKAKPSSL LPACLKMLQ Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.
Characteristics:	Key Benefits: • Made in Germany - from design to production - by highly experienced protein experts. • Protein expressed with ALiCE® and purified in one-step affinity chromatography • These proteins are normally active (enzymatically functional) as our customers have

- reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	CIDEC
Alternative Name:	Cidec (CIDEC Products)
Background:	Lipid transferase CIDEC (Cell death-inducing DFFA-like effector protein C) (Fat-specific protein

Target Details

FSP27),FUNCTION: Lipid transferase specifically expressed in white adipose tissue, which promotes unilocular lipid droplet formation by mediating lipid droplet fusion (PubMed:18334488, PubMed:22144693, PubMed:26733203, PubMed:30361435, PubMed:36477540). Lipid droplet fusion promotes their enlargement, restricting lipolysis and favoring lipid storage (PubMed:18334488, PubMed:18682832, PubMed:22144693, PubMed:26733203). Localizes on the lipid droplet surface, at focal contact sites between lipid droplets, and mediates atypical lipid droplet fusion by undergoing liquid-liquid phase separation (LLPS) and promoting directional net neutral lipid transfer from the smaller to larger lipid droplets (PubMed:18334488, PubMed:22144693). The transfer direction may be driven by the internal pressure difference between the contacting lipid droplet pair (PubMed:18334488, PubMed:22144693). Its role in neutral lipid transfer and lipid droplet enlargement is activated by the interaction with PLIN1 (PubMed:23481402). May also act as a CEBPB coactivator in the white adipose tissue to control the expression of a subset of CEBPB downstream target genes, including SOCS1, SOCS3, TGFB1, TGFB1, ID2 and XDH (PubMed:22245780). When overexpressed in preadipocytes, induces apoptosis or increases cell susceptibility to apoptosis induced by serum deprivation or TGFB treatment (By similarity). {ECO:0000250|UniProtKB:Q96AQ7, ECO:0000269|PubMed:18334488, ECO:0000269|PubMed:18682832, ECO:0000269|PubMed:22144693, ECO:0000269|PubMed:22245780, ECO:0000269|PubMed:23481402, ECO:0000269|PubMed:26733203, ECO:0000269|PubMed:30361435, ECO:0000269|PubMed:36477540}.

Molecular Weight: 27.3 kDa

UniProt: [P56198](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the

Application Details

mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.
Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months