

Datasheet for ABIN3134755

CD130/gp130 Protein (AA 640-917) (His tag)[Go to Product page](#)**1** Image

Overview

| | |
|-------------------------------|--|
| Quantity: | 1 mg |
| Target: | CD130/gp130 (IL6ST) |
| Protein Characteristics: | AA 640-917 |
| Origin: | Mouse |
| Source: | Escherichia coli (E. coli) |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CD130/gp130 protein is labelled with His tag. |
| Application: | SDS-PAGE (SDS), Western Blotting (WB), ELISA, Crystallization (Crys) |

Product Details

| | |
|------------------|--|
| Sequence: | <p>NKRDLIKHHI WPNVPDPSKS HIAQWSPHTP PRHNFNSKDQ MYSDGNFTDV SVVEIANNK KPCPDDLKSV DLFKKEKVST EGHSSGIGGS SCMSSSRPSI SSNEENESAQ STASTVQYST VHSGYRHHV PSVQVFSRSE STQPLLDSEE RPEDLQLVDS VGGDEILPR QPYFKQNCST PEACPEISHF ERSNQVLSGN EEDFVRLKQQ QVSDHISQPY GSEQRRLFQE GSTADALGTG ADGQMERFES VGMETTIDEE IPKSYLPQTV RQGGYMPQ</p> <p>Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.</p> |
| Characteristics: | <ul style="list-style-type: none">• Made in Germany - from design to production - by highly experienced protein experts.• Mouse Il6st Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.• State-of-the-art algorithm used for plasmid design (Gene synthesis). <p>This protein is a made to order protein and will be made for the first time for your order. Our</p> |

Product Details

experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

| | |
|------------------|---|
| Purification: | Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot. |
| Purity: | >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | Endotoxin has not been removed. Please contact us if you require endotoxin removal. |
| Grade: | Crystallography grade |

Target Details

| | |
|-------------------|---|
| Target: | CD130/gp130 (IL6ST) |
| Alternative Name: | Il6st (IL6ST Products) |
| Background: | Signal-transducing molecule. The receptor systems for IL6, LIF, OSM, CNTF, IL11, CTF1 and BSF3 can utilize IL6ST for initiating signal transmission. Binding of IL6 to IL6R induces IL6ST |

Target Details

homodimerization and formation of a high-affinity receptor complex, which activates Janus kinases (PubMed:1602143). That causes phosphorylation of IL6ST tyrosine residues which in turn activates STAT3 (PubMed:10661409). Mediates signals which regulate immune response, hematopoiesis, pain control and bone metabolism (PubMed:10661409, PubMed:26255596, PubMed:25057188, PubMed:8552649). Has a role in embryonic development (PubMed:10661409). Does not bind IL6 (By similarity). Essential for survival of motor and sensory neurons and for differentiation of astrocytes (PubMed:10377352). Required for expression of TRPA1 in nociceptive neurons (PubMed:25057188). Required for the maintenance of PTH1R expression in the osteoblast lineage and for the stimulation of PTH-induced osteoblast differentiation (PubMed:25228504). Required for normal trabecular bone mass and cortical bone composition (PubMed:24339143, PubMed:9348227, PubMed:26255596). {ECO:0000250|UniProtKB:P40189, ECO:0000269|PubMed:10377352, ECO:0000269|PubMed:10661409, ECO:0000269|PubMed:1602143, ECO:0000269|PubMed:24339143, ECO:0000269|PubMed:25057188, ECO:0000269|PubMed:25228504, ECO:0000269|PubMed:26255596, ECO:0000269|PubMed:8552649, ECO:0000269|PubMed:9348227}.

Molecular Weight: 32.0 kDa Including tag.

UniProt: [Q00560](#)

Pathways: [JAK-STAT Signaling](#), [Cellular Glucan Metabolic Process](#), [Autophagy](#), [Smooth Muscle Cell Migration](#), [Cancer Immune Checkpoints](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

Handling

Format: Liquid

Handling

| | |
|------------------|--|
| Buffer: | 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. |
| Expiry Date: | Unlimited (if stored properly) |

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process