

Datasheet for ABIN3134820

ITK Protein (AA 1-625) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	ITK
Protein Characteristics:	AA 1-625
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This ITK protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Brand:	AliCE®
Sequence:	<p>MNNFILLEEQ LIKKSQQKRR TSPSNFKVRF FVLTKASLAY FEDRHGKKRT LKGSIELSRI</p> <p>KCVEIVKSDI SIPCHYKYPF QTLVYLQVVH DNYLLYVFAP DCESRQRWVL TLKEETRNNN</p> <p>SLVSKYHPNF WMDGRWRCCS QLEKPAVGCA PYDPSKNASK KPLPPTPEDN RRSFQEPEET</p> <p>LVIALYDYQT NDPQELALRC DEEYLLDSS EIHWWRVQDK NGHEGYAPSS YLVEKSPNNL</p> <p>ETYEWYNKSI SRDKAEKLLL DTGKEGAFMV RDSRTPGTYT VSVFTKAIIS ENPCIKHYHI</p> <p>KETNDSPKRY YVAEKYVFDS IPLLQYHQY NGGGLVTRLR YPVCWRQKA PVTAGLRYGK</p> <p>WVIQPSLTF VQEIGSGQFG LVHLGYWLNK DKVAIKTIQE GAMSEEDFIE EAEVMMKLSH</p> <p>PKLVQLYGVC LEQAPICLVF EFMEHGCLSD YLRSQRGLFA AETLLGMCLD VCEGMAYLEK</p> <p>ACVIHRDLAA RNCLVGENQV IKVSDFGMTR FVLDDQYTSS TGTKFPVKWA SPEVFSFSRY</p> <p>SSKSDVWSFG VLMWEVFSEG KIPYENRSNS EVVEDISTGF RLYKPRLASC HVYQIMNHCW</p> <p>KEKPEDRPPF SQLLSQLAEI AEAGL</p>

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Product Details

Grade: custom-made

Target Details

Target: ITK

Alternative Name: Itk ([ITK Products](#))

Background: Tyrosine-protein kinase ITK/TSK (EC 2.7.10.2) (IL-2-inducible T-cell kinase) (Kinase EMT) (Kinase TLK) (T-cell-specific kinase),FUNCTION: Tyrosine kinase that plays an essential role in regulation of the adaptive immune response. Regulates the development, function and differentiation of conventional T-cells and nonconventional NKT-cells. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylation lead to the recruitment of ITK to the cell membrane, in the vicinity of the stimulated TCR receptor, where it is phosphorylated by LCK. Phosphorylation leads to ITK autophosphorylation and full activation. Once activated, phosphorylates PLCG1, leading to the activation of this lipase and subsequent cleavage of its substrates. In turn, the endoplasmic reticulum releases calcium in the cytoplasm and the nuclear activator of activated T-cells (NFAT) translocates into the nucleus to perform its transcriptional duty. Phosphorylates 2 essential adapter proteins: the linker for activation of T-cells/LAT protein and LCP2. Then, a large number of signaling molecules such as VAV1 are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation. Required for TCR-mediated calcium response in gamma-delta T-cells, may also be involved in the modulation of the transcriptomic signature in the Vgamma2-positive subset of immature gamma-delta T-cells (PubMed:23562159). Phosphorylates TBX21 at 'Tyr-525' and mediates its interaction with GATA3 (PubMed:15662016). {ECO:0000269|PubMed:15662016, ECO:0000269|PubMed:21036902, ECO:0000269|PubMed:23562159}.

Molecular Weight: 72.3 kDa

UniProt: [Q03526](#)

Pathways: [TCR Signaling, Fc-epsilon Receptor Signaling Pathway](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce

Application Details

even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months