antibodies .- online.com





ECT2 Protein (AA 2-913) (His tag)



Overview

Quantity:	1 mg
Target:	ECT2
Protein Characteristics:	AA 2-913
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ECT2 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:

ADDSVLPSPS EITSLADSSV FDSKVAEMSK ENLCLASTSN VDEEMPQVEA RVIMVQDAGK
QEELLKALKT IKIMEVPVIK IKESCPGKSE EKLIKSIINM EMKVPCVKMD SMEEFESLDS
PEFENIFVVT DFQNSVFNDL YKADCRIVGP PVILNCAQRG EPLPFSCRPL YCTSMLNLVL
CFTGFRKKEE LVKLVTLVHH MGGVIRKECN SKVTHLVANC TQGEKFRVAV SLGTPIMKPE
WIYKAWERRN EQCFCAAVDD FRNEFKVPPF QDCILSFLGF SDEEKHSMEE MTEMQGGSYL
PVGDERCTHL IVEENTVKDL PFEPSKKLFV VKQEWFWGSI QMDARAGETM YLYEKANTPE
LKKSVSLLSL STPNSNRKRR RLKETLAQLS RETDLSPFPP RKRPSAEHSL SIGSLLDISN
TPESSIHYGE TPKSCAKSSR SSTPVPPKQS ARWQVAKELY QTESNYVNIL ATIIQLFQVP
LEEEGQRGGP ILAPEEIKTI FGSIPDIFDV HMKIKDDLED LIANWDESRS IGDIFLKYAK
DLVKTYPPFV NFFEMSKEMI IKCEKQKPRF HAFLKINQAK PECGRQSLVE LLIRPVQRLP
SVALLLNDLK KHTADENPDK STLEKAIGSL KEVMTHINED KRKTEAQKQI FDVVYEVDGC
PANLLSSHRS LVQRVETVSL GEHPCDRGEQ VTLFLFNDCL EIARKRHKVI GTFRSPHDRT

RPPASLKHIH LMPLSQIKKV LDIRETEDCH NAFALLVRPP TEQANVLLSF QMTSEELPKE SWLKMLCRHV ANTICKADAE NLMYVADPES FEVNTKDMDS TLSRASRAIK KTSKKVTRAF SFSKTPKRAL RMALSSSHSS EGRSPPSSGK LAVSRLSSTS SLAGIPSPSL VSLPSFFERR SHTLSRSTTH LI

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Ect2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Product Details Sterility: 0.22 µm filtered Endotoxin Level: Protein is endotoxin free. Grade: Crystallography grade **Target Details** ECT2 Target: Alternative Name: Ect2 (ECT2 Products) Guanine nucleotide exchange factor (GEF) that catalyzes the exchange of GDP for GTP. Background: Promotes guanine nucleotide exchange on the Rho family members of small GTPases, like RHOA, RHOC, RAC1 and CDC42. Required for signal transduction pathways involved in the regulation of cytokinesis. Component of the centralspindlin complex that serves as a microtubule-dependent and Rho-mediated signaling required for the myosin contractile ring formation during the cell cycle cytokinesis. Regulates the translocation of RHOA from the central spindle to the equatorial region. Plays a role in the control of mitotic spindle assembly, regulates the activation of CDC42 in metaphase for the process of spindle fibers attachment to kinetochores before chromosome congression. Involved in the regulation of epithelial cell polarity, participates in the formation of epithelial tight junctions in a polarity complex PARD3-PARD6-protein kinase PRKCQ-dependent manner. Plays a role in the regulation of neurite outgrowth. Inhibits phenobarbital (PB)-induced NR1I3 nuclear translocation. Stimulates the activity of RAC1 through its association with the oncogenic PARD6A-PRKCI complex in cancer cells, thereby acting to coordinately drive tumor cell proliferation and invasion. Also stimulates genotoxic stress-induced RHOB activity in breast cancer cells leading to their cell death. {ECO:0000269|PubMed:17904126, ECO:0000269|PubMed:21350944}. Molecular Weight: 104.0 kDa Including tag. UniProt: Q07139 Pathways: Neurotrophin Signaling Pathway, Cell-Cell Junction Organization **Application Details** In addition to the applications listed above we expect the protein to work for functional studies **Application Notes:** as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.

Protein has not been tested for activity yet. In cases in which it is highly likely that the

Comment:

Application Details

	recombinant protein with the default tag will be insoluble our protein lab may suggest a higher
	molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible
	options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)