

Datasheet for ABIN3134854

MAP3K8 Protein (AA 1-467) (His tag)**1** Image[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	MAP3K8
Protein Characteristics:	AA 1-467
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MAP3K8 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)

Product Details

Sequence:	MEYMSTGSDE KEEIDLLIKH LNVSEVIDIM ENLYASEEPG VYEPSLMTMY PDSNQNEERS ESLLRSGQEV PWLSSVRYGT VEDLLAFANH VSNMTKHFYR RRPQECGILL NMVISPQNGR YQIDSDVLLV PWKLTyrNIG SGFVPRGAFG KVYLAQDMKT KKRmACKLIP IDQFKPSDVE IQACFRHENI AELYGAVLWG DTVHLFMEAG EGGSVLEKLE SCGPMREFEI IWVTKHILKG LDFLHsKKVI HHDIKPSNIV FMSTKAVLVD FGLSVKMTED VYLPKDLRGT EIYMSPEVIL CRGHSTKADI YSLGATLIHM QTGTppWVKR YPRsAYPSYL YIIHKQAPPL EDIAGDCSPG MRELIEAALE RNPnHRPKAA DLLKHEALNP PREDQPRCQS LDSALFERKR LLSRKELQLP ENIADSSCTG STEESEVLRR QRSlyIDLGA LAGYFNIVRG PPTLEYG Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.
Characteristics:	<ul style="list-style-type: none">• Made in Germany - from design to production - by highly experienced protein experts.• Mouse Map3k8 Protein (raised in Insect Cells) purified by multi-step, protein-specific process

to ensure crystallization grade.

- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the ExPASy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

Target:

MAP3K8

Alternative Name:

Map3k8 ([MAP3K8 Products](#))

Target Details

Background:	Required for lipopolysaccharide (LPS)-induced, TLR4-mediated activation of the MAPK/ERK pathway in macrophages, thus being critical for production of the proinflammatory cytokine TNF-alpha (TNF) during immune responses. Involved in the regulation of T-helper cell differentiation and IFNG expression in T-cells. Involved in mediating host resistance to bacterial infection through negative regulation of type I interferon (IFN) production. In vitro, activates MAPK/ERK pathway in response to IL1 in an IRAK1-independent manner, leading to up-regulation of IL8 and CCL4. Transduces CD40 and TNFRSF1A signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation of immunoglobulin production. May also play a role in the transduction of TNF signals that activate JNK and NF-kappa-B in some cell types. In adipocytes, activates MAPK/ERK pathway in an IKBKB-dependent manner in response to IL1B and TNF, but not insulin, leading to induction of lipolysis. Plays a role in the cell cycle. {ECO:0000269 PubMed:11163183, ECO:0000269 PubMed:12667451, ECO:0000269 PubMed:12881420, ECO:0000269 PubMed:15833743, ECO:0000269 PubMed:19001140, ECO:0000269 PubMed:19808894, ECO:0000269 PubMed:23842752}.
Molecular Weight:	53.9 kDa Including tag.
UniProt:	Q07174
Pathways:	PI3K-Akt Signaling , TCR Signaling

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling

Handling Advice:	Avoid repeated freeze-thaw cycles.
------------------	------------------------------------

Storage:	-80 °C
----------	--------

Storage Comment:	Store at -80°C.
------------------	-----------------

Expiry Date:	Unlimited (if stored properly)
--------------	--------------------------------

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process