antibodies .- online.com





PPP4R2 Protein (AA 1-417) (His tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	PPP4R2
Protein Characteristics:	AA 1-417
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PPP4R2 protein is labelled with His tag.
Application:	ELISA, Crystallization (Crys), SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Sequence:

MDVERLQEAL KDFEKRGKKE VCPVLDQFLC HVAKTGETMI QWSQFKGYFI FKLEKVMDDF
RTSAPEPRGP PNPNVEYIPF DEMKERILKI VTGFNGIPFT IQRLCELLTD PRRNYTGTDK
FLRGVEKNVM VVSCVCPSSE KNNSNSLNRM NGVMFPGNSP NYTDRSNING PGTPRPLNRP
KLSLSAPLTT NGLPESTDSK DSELQLSEEK GHSDSSASES EVSLLSPVKN KHPDEDAVES
EEHEVKRLKF DKEGDVRETA SQTVSGEVSS VRAEETETAA PPPDKDRESR TRQHCTEEEE
EEEEEEEEEE EESFMTPREM VPERKNQEKE SDDALTVNEE TSEESHQMEG SGVSPAQTDS
TSERSDSAGA SRSGSDCLET QESGGPPSSK TGESVSVPSS MESEEATEVT DDPMEQD

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Ppp4r2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 μm filtered
Endotoxin Level:	Protein is endotoxin free.
Grade:	Crystallography grade

Target Details

Target:	PPP4R2
Alternative Name:	Ppp4r2 (PPP4R2 Products)

Target Details

Background:	Regulatory subunit of serine/threonine-protein phosphatase 4 (PP4). May regulate the activity
	of PPP4C at centrosomal microtubule organizing centers. Its interaction with the SMN complex
	leads to enhance the temporal localization of snRNPs, suggesting a role of PPP4C in
	maturation of spliceosomal snRNPs. The PPP4C-PPP4R2-PPP4R3A PP4 complex specifically
	dephosphorylates H2AFX phosphorylated on 'Ser-140' (gamma-H2AFX) generated during DNA
	replication and required for DNA double strand break repair (By similarity). Mediates RPA2
	dephosphorylation by recruiting PPP4C to RPA2 in a DNA damage-dependent manner. RPA2
	dephosphorylation is required for the efficient RPA2-mediated recruitment of RAD51 to
	chromatin following double strand breaks, an essential step for DNA repair (By similarity).
	{ECO:0000250}.
Molecular Weight:	47.4 kDa Including tag.
UniProt:	Q0VGB7
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the
	recombinant protein with the default tag will be insoluble our protein lab may suggest a higher
	molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible
	options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process