antibodies -online.com





ATP8B1 Protein (AA 1-121) (His tag)



Go to Product page

\sim	
()\/△	rview
\circ	1 410 44

Quantity:	1 mg
Target:	ATP8B1
Protein Characteristics:	AA 1-121
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This ATP8B1 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)
Product Details	
Sequence:	MSTERDSETT FDEESQPNDE VVPYSDDETE DELEDQGSTV EPEQNRVNRE AEKKRETFRK
	DCTWQVKAND RKFHEQPHFM NTKFFCIKES KYASNAIKTY KYNGFTFLPM NLFEQFKRAA N
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	 Made in Germany - from design to production - by highly experienced protein experts. Mouse Atp8b1 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made to order protein and will be made for the first time for your order. Our
	experts in the lab will ensure that you receive a correctly folded protein.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom
	made proteins from other companies is that there is no financial obligation in case the protein

cannot be expressed or purified.

specific reference buffer.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

Target Details

Target:	ATP8B1
Alternative Name:	Atp8b1 (ATP8B1 Products)
Background:	Catalytic component of a P4-ATPase flippase complex which catalyzes the hydrolysis of ATP
	coupled to the transport of aminophospholipids from the outer to the inner leaflet of various
	membranes and ensures the maintenance of asymmetric distribution of phospholipids.
	Phospholipid translocation seems also to be implicated in vesicle formation and in uptake of
	lipid signaling molecules. May play a role in asymmetric distribution of phospholipids in the

canicular membrane. Plays a role in bile salt homeostasis. In cooperation with ABCB4 may be involved in establishing integrity of the canalicular membrane thus protecting hepatocytes from bile salts. Involved in the microvillus formation in polarized epithelial cells, the function seems to be independent from its flippase activity. Required for the preservation of cochlear hair cells in the inner ear. Required for the preservation of cochlear hair cells in the inner ear. According PubMed:20852622 is proposed to act as cardiolipin transporter during inflammatory injury, the function is questioned by PubMed:21475228. {ECO:0000269|PubMed:14976163, ECO:0000269|PubMed:19478059, ECO:0000269|PubMed:21820390}.

Molecular Weight:

15.4 kDa Including tag.

UniProt:

Q148W0

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.

Comment:

Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)