antibodies .- online.com





CDT2/RAMP Protein (AA 1-729) (Strep Tag)



Go to Product page

Overview

Quantity:	1 mg
Target:	CDT2/RAMP (DTL)
Protein Characteristics:	AA 1-729
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CDT2/RAMP protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Sequence:

MLFNSVLRQP QLGVLRNGWS SHYPLQSLLS GYQCNCNDEH TSYGETGVPV PPFGCTFCTA
PSMEHILAVA NEEGFVRLYN TESQTSKKTC FKEWMAHWNA VFDLAWVPGE LKLVTAAGDQ
TAKFWDVRAG ELMGTCKGHQ CSLKSVAFPK FQKAVFSTGG RDGNIMIWDT RCNKKDGFYR
QVNQISGAHN TADKQTPSKP KKKQNSKGLA PAVDSQQSVT VVLFQDENTL VSAGAVDGII
KVWDLRKNYT AYRQEPIASK SFLYPGTSTR KLGYSSLVLD STGSTLFANC TDDNIYMFNM
TGLKTSPVAV FNGHQNSTFY VKSSLSPDDQ FLISGSSDEA AYIWKVSMPW HPPTVLLGHS
QEVTSVCWCP SDFTKIATCS DDNTLKIWRL NRGLEEKPGD KHSIVGWTSQ KKKEVKACPV
TVPSSQSTPA KAPRAKSSPS ISSPSSAACT PSCAGDLPLP SSTPTFSVKT TPATTRSSVS
RRGSISSVSP KPLSSFKMSL RNWVTRTPSS SPPVTPPASE TKISSPRKAL IPVSQKSSQA
DACSESRNRV KRRLDSSCLE SVKQKCVKSC NCVTELDGQA ESLRLDLCCL SGTQEVLSQD
SEGPTKSSKT EGAGTSISEP PSPVSPYASE GCGPLPLPLR PCGEGSEMVG KENSSPENKN
WLLAIAAKRK AENSSPRSPS SQTPSSRRQS GKTSPGPVTI TPSSMRKICT YFRRKTQDDF

CSPEHSTEL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- · The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System

(ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level:

Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Grade:

Crystallography grade

Target Details

Target:

CDT2/RAMP (DTL)

Alternative Name:

Dtl (DTL Products)

Background:

Denticleless protein homolog (Lethal(2) denticleless protein homolog) (Meth A retinoic acidregulated nuclear matrix-associated protein) (Meth A RAMP) (Retinoic acid-regulated nuclear matrix-associated protein),FUNCTION: Substrate-specific adapter of a DCX (DDB1-CUL4-X-box) E3 ubiquitin-protein ligase complex required for cell cycle control, DNA damage response and translesion DNA synthesis. The DCX(DTL) complex, also named CRL4(CDT2) complex, mediates the polyubiquitination and subsequent degradation of CDT1, CDKN1A/p21(CIP1), FBH1, KMT5A and SDE2. CDT1 degradation in response to DNA damage is necessary to ensure proper cell cycle regulation of DNA replication. CDKN1A/p21(CIP1) degradation during S phase or following UV irradiation is essential to control replication licensing. KMT5A degradation is also important for a proper regulation of mechanisms such as TGF-beta signaling, cell cycle progression, DNA repair and cell migration. Most substrates require their interaction with PCNA for their polyubiquitination: substrates interact with PCNA via their PIP-box, and those containing the 'K+4' motif in the PIP box, recruit the DCX(DTL) complex, leading to their degradation. In undamaged proliferating cells, the DCX(DTL) complex also promotes the 'Lys-164' monoubiquitination of PCNA, thereby being involved in PCNA-dependent translesion DNA synthesis. The DDB1-CUL4A-DTL E3 ligase complex regulates the circadian clock function by mediating the ubiquitination and degradation of CRY1 (By similarity). {ECO:0000250|UniProtKB:Q9NZJ0}.

Molecular Weight:

79.1 kDa

UniProt:

O3TLR7

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)