antibodies .- online.com





THRAP3 Protein (AA 2-951) (His tag)



Overview

Quantity:	1 mg
Target:	THRAP3
Protein Characteristics:	AA 2-951
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This THRAP3 protein is labelled with His tag.
Application:	Crystallization (Crys), ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Sequence:

SKTNKSKSGS RSSRSRSASR SRSRSFSKSR SRSRSVSRSR KRRLSSRSRS RSYSPAHNRE
RNHPRVYQNR DFRGHNRGYR RPYYFRGRNR GFYPWGQYNR GGYGNYRSNW QNYRQAYSPR
RGRSRSRSPK RRSPSPRSRS HSRNSDKSSS DRSRRSSSSR SSSNHSRVES SKRKSTKEKK
SSSKDSRPSQ AAGDNQGDEA KEQTFSGGTS QDIKGSESSK PWPDATTYGA GSASRASVSD
LSPRERSPAL KSPLQSVVVR RRSPRPSPVP KPSPPLSNAS QMGSSMSGGA GYQSGAHQGQ
FDHGSGSLSP SKKSPVGKSP PATGSAYGSS QKEESAASGG AAYSKRYLEE QKTENGKDKE
QKQTNADKEK LKEKGGFSDA DVKMKSDPFA PKTDSEKPFR GSQSPKRYKL RDDFEKKMAD
FHKEELDEHD KDKSKGRKEP EFDDEPKFMS KVIAGASKNQ EEEKSGKWES LHTGKEKQRK
AEEMEDEPFT ERSRKEERGG SKRSESGHRG FVPEKNFRVT AYKAVQEKSS SPPPRKTSES
RDKLGSKGDF SSGKSSFSIT REAQVNVRMD SFDEDLARPS GLLAQERKLC RDLVHSNKKE
QEFRSIFQHI QSAQSQRSPS ELFAQHIVTI VHHVKEHHFG SSGMTLHERF TKYLKRGNEQ
EAAKNKKSPE IHRRIDISPS TFRKHGLTHE ELKSPREPGY KAEGKYKDDP VDLRLDIERR

KKHKERDLKR GKSRESVDSR DSSHSRERST EKTEKTHKGS KKQKKHRRAR DRSRSSSSS QSSHSYKAEE YPEEAEEREE STSGFDKSRL GTKDFVGPNE RGGRARGTFQ FRARGRGWGR GNYSGNNNNN SNNDFQKRSR EEEWDPEYTP KSKKYYLHDD REGEGSDKWM GRGRGRGAFP RGRGRFMFRK SSTSPKWAHD KFSGEEGEIE DDESGTENRE EKDSLQPSAE

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Thrap3 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Product Details Sterility: 0.22 µm filtered Endotoxin Level: Protein is endotoxin free. Grade: Crystallography grade **Target Details** THRAP3 Target: Alternative Name: Thrap3 (THRAP3 Products) Background: Involved in pre-mRNA splicing. Remains associated with spliced mRNA after splicing which probably involves interactions with the exon junction complex (EJC). Can trigger mRNA decay which seems to be independent of nonsense-mediated decay involving premature stop codons (PTC) recognition. May be involved in nuclear mRNA decay. Involved in regulation of signalinduced alternative splicing. During splicing of PTPRC/CD45 is proposed to sequester phosphorylated SFPQ from PTPRC/CD45 pre-mRNA in resting T-cells. Involved in cyclin-D1/CCND1 mRNA stability probably by acting as component of the SNARP complex which associates with both the 3'end of the CCND1 gene and its mRNA. Involved in response to DNA damage. Is excluded from DNA damage sites in a manner that parallels transcription inhibition, the function may involve the SNARP complex. Initially thought to play a role in transcriptional coactivation through its association with the TRAP complex, however, it is not regarded as a stable Mediator complex subunit. Cooperatively with HELZ2, enhances the transcriptional activation mediated by PPARG, maybe through the stabilization of the PPARG binding to DNA in presence of ligand. May play a role in the terminal stage of adipocyte differentiation. Plays a role in the positive regulation of the circadian clock. Acts as a coactivator of the CLOCK-ARNTL/BMAL1 heterodimer and promotes its transcriptional activator activity and binding to circadian target genes (PubMed:24043798). {ECO:0000269|PubMed:23525231, ECO:0000269|PubMed:24043798}. Molecular Weight:

Molecular Weight: 109.0 kDa Including tag.

UniProt: Q569Z6

Pathways: Intracellular Steroid Hormone Receptor Signaling Pathway

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee

Application Details

	though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)