
antibodies .- online.com





FBXW11 Protein (AA 1-542) (Strep Tag)



Go to Product page

Overview

Quantity:	1 mg
Target:	FBXW11
Protein Characteristics:	AA 1-542
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This FBXW11 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence:

MEPDSVIEDK TIELMCSVPR SLWLGCANLV ESMCALSCLQ SMPSVRCLQI SNGTSSVIVS
RKRPSEGNYQ KEKDLCIKYF DQWSESDQVE FVEHLISRMC HYQHGHINSY LKPMLQRDFI
TALPEQGLDH IAENILSYLD ARSLCAAELV CKEWQRVISE GMLWKKLIER MVRTDPLWKG
LSERRGWDQY LFKNRPTDGP PNSFYRSLYP KIIQDIETIE SNWRCGRHNL QRIQCRSENS
KGVYCLQYDD DKIISGLRDN SIKIWDKSSL ECLKVLTGHT GSVLCLQYDE RVIVTGSSDS
TVRVWDVNTG EVLNTLIHHN EAVLHLRFSN GLMVTCSKDR SIAVWDMASA TDITLRRVLV
GHRAAVNVVD FDDKYIVSAS GDRTIKVWST STCEFVRTLN GHKRGIACLQ YRDRLVVSGS
SDNTIRLWDI ECGACLRVLE GHEELVRCIR FDNKRIVSGA YDGKIKVWDL QAALDPRAPA
STLCLRTLVE HSGRVFRLQF DEFQIISSSH DDTILIWDFL NVPPSAQNET RSPSRTYTYI SR

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and

Product Details		
	Western blot.	
Purity:	≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.	
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)	
Target Details		
Target:	FBXW11	
Alternative Name:	Fbxw11 (FBXW11 Products)	
Background:	F-box/WD repeat-containing protein 11 (F-box and WD repeats protein beta-TrCP2) (F-box/WD repeat-containing protein 1B) (Homologous to Slimb protein) (HOS), FUNCTION: Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:11896578). Probably recognizes and binds to phosphorylated target proteins: the interaction with substrates requires the phosphorylation of the two serine residues in the substrates' destruction motif D-S-G-X(2,3,4)-S (By similarity). SCF(FBXW11) mediates the ubiquitination of phosphorylated CTNNB1 and participates in Wnt signaling regulation (By similarity). SCF(FBXW11) plays a key role in NF-kappa-B activation by mediating ubiquitination of phosphorylated NFKBIA, leading to its degradation by the proteasome, thereby allowing the associated NF-kappa-B complex to translocate into the nucleus and to activate transcription (PubMed:11896578). The SCF(FBXW11) complex also regulates NF-kappa-B by mediating ubiquitination of phosphorylated NFKB1: specifically ubiquitinates the p105 form of NFKB1, leading to its degradation (By similarity). SCF(FBXW11) mediates the ubiquitination of IFNAR1 (By similarity). SCF(FBXW11) mediates the ubiquitination of CEP68, this is required for centriole separation during mitosis (By similarity). Involved in the oxidative stress-induced a ubiquitin-mediated decrease in RCAN1 (By similarity). Mediates the degradation of CDC25A induced by ionizing radiation in cells progressing through S phase and thus may function in the intra-S-phase checkpoint (By similarity). Has an essential role in the control of the clock-dependent transcription via degradation of phosphorylated PER1 and phosphorylated PER2 (PubMed:18782782). SCF(FBXW11) mediates the ubiquitination of CYTH1, and probably CYTH2 (By similarity). SCF(FBXW11) acts as a regulator of mTORC1 signaling pathway by catalyzing ubiquitination and subsequent proteasomal degradation	

Molecular Weight:

62.1 kDa

Target Details UniProt: Q5SRY7 **Application Details** In addition to the applications listed above we expect the protein to work for functional studies Application Notes: as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though. ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Comment: Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein! Restrictions: For Research Use only Handling Format: Liquid Buffer: The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.

Avoid repeated freeze-thaw cycles.

Unlimited (if stored properly)

-80 °C

Store at -80°C.

Handling Advice:

Storage Comment:

Storage:

Expiry Date: