antibodies .- online.com





PP5 Protein (AA 2-499) (His tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	PP5 (PPP5C)
Protein Characteristics:	AA 2-499
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PP5 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

Product Details

Sequence:

AMAEGERTEC AETPRDEPPA DGTLKRAEEL KTQANDYFKA KDYENAIKFY SQAIELNPGN
AIYYGNRSLA YLRTECYGYA LGDATRAIEL DKKYIKGYYR RAASNMALGK FRAALRDYET
VVKVKPNDKD AKMKYQECSK IVKQKAFERA IAGDEHRRSV VDSLDIESMT IEDEYSGPKL
EDGKVTITFM KDLMQWYKDQ KKLHRKCAYQ ILVQVKEVLC KLSTLVETTL KETEKITVCG
DTHGQFYDLL NIFELNGLPS ETNPYIFNGD FVDRGSFSVE VILTLFGFKL LYPDHFHLLR
GNHETDNMNQ IYGFEGEVKA KYTAQMYELF SEVFEWLPLA QCINGKVLIM HGGLFSEDGV
TLDDIRKIER NRQPPDSGPM CDLLWSDPQP QNGRSVSKRG VSCQFGPDVT KAFLEENQLD
YIIRSHEVKA EGYEVAHGGR CVTVFSAPNY CDQMGNKASY IHLQGSDLRP QFHQFTAVPH
PNVKPMAYAN TLLQLGMM

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Ppp5c Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

Target:	PP5 (PPP5C)
Alternative Name:	Ppp5c (PPP5C Products)
Background:	Serine/threonine-protein phosphatase that dephosphorylates a myriad of proteins involved in
	different signaling pathways including the kinases CSNK1E, ASK1/MAP3K5, PRKDC and RAF1,
	the nuclear receptors NR3C1, PPARG, ESR1 and ESR2, SMAD proteins and TAU/MAPT.
	Implicated in wide ranging cellular processes, including apoptosis, differentiation, DNA damage
	response, cell survival, regulation of ion channels or circadian rhythms, in response to steroid
	and thyroid hormones, calcium, fatty acids, TGF-beta as well as oxidative and genotoxic
	stresses. Participates in the control of DNA damage response mechanisms such as checkpoint
	activation and DNA damage repair through, for instance, the regulation ATM/ATR-signaling and
	dephosphorylation of PRKDC and TP53BP1. Inhibits ASK1/MAP3K5-mediated apoptosis
	induced by oxidative stress. Plays a positive role in adipogenesis, mainly through the
	dephosphorylation and activation of PPARG transactivation function. Also dephosphorylates
	and inhibits the anti-adipogenic effect of NR3C1. Regulates the circadian rhythms, through the
	dephosphorylation and activation of CSNK1E. May modulate TGF-beta signaling pathway by
	the regulation of SMAD3 phosphorylation and protein expression levels. Dephosphorylates and
	may play a role in the regulation of TAU/MAPT. Through their dephosphorylation, may play a
	role in the regulation of ions channels such as KCNH2. {ECO:0000269 PubMed:17376776,
	ECO:0000269 PubMed:21994940, ECO:0000269 PubMed:22526606}.
Molecular Weight:	57.7 kDa Including tag.
UniProt:	Q60676
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the
	recombinant protein with the default tag will be insoluble our protein lab may suggest a higher
	molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible
	options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process