antibodies .- online.com





MST1R Protein (AA 25-960) (His tag)



Image



Overview

Quantity:	1 mg
Target:	MST1R
Protein Characteristics:	AA 25-960
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MST1R protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:

TNLNWQCPRI PYAASRDFSV KYVVPSFSAG GRVQATAAYE DSTNSAVFVA TRNHLHVLGP
DLQFIENLTT GPIGNPGCQT CASCGPGPHG PPKDTDTLVL VMEPGLPALV SCGSTLQGRC
FLHELEPRGK ALHLAAPACL FSANNNKPEA CTDCVASPLG TRVTVVEQGH ASYFYVASSL
DPELAASFSP RSVSIRRLKS DTSGFQPGFP SLSVLPKYLA SYLIKYVYSF HSGDFVYFLT
VQPISVTSPP SALHTRLVRL NAVEPEIGDY RELVLDCHFA PKRRRRGAPE GTQPYPVLQA
AHSAPVDAKL AVELSISEGQ EVLFGVFVTV KDGGSGMGPN SVVCAFPIYH LNILIEEGVE
YCCHSSNSSS LLSRGLDFFQ TPSFCPNPPG GEASGPSSRC HYFPLMVHAS FTRVDLFNGL
LGSVKVTALH VTRLGNVTVA HMGTVDGRVL QVEIARSLNY LLYVSNFSLG SSGQPVHRDV
SRLGNDLLFA SGDQVFKVPI QGPGCRHFLT CWRCLRAQRF MGCGWCGDRC DRQKECPGSW
QQDHCPPEIS EFYPHSGPLR GTTRLTLCGS NFYLRPDDVV PEGTHQITVG QSPCRLLPKD
SSSPRPGSLK EFIQELECEL EPLVTQAVGT TNISLVITNM PAGKHFRVEG ISVQEGFSFV
EPVLTSIKPD FGPRAGGTYL TLEGQSLSVG TSRAVLVNGT QCRLEQVNEE QILCVTPPGA

GTARVPLHLQ IGGAEVPGSW TFHYKEDPIV LDISPKCGYS GSHIMIHGQH LTSAWHFTLS
FHDGQSTVES RCAGQFVEQQ QRRCRLPEYV VRNPQGWATG NLSVWGDGAA GFTLPGFRFL
PPPSPLRAGL VELKPEEHSV KVEYVGLGAV ADCVTVNMTV GGEVCQHELR GDVVICPLPP
SLQLGKDGVP LQVCVDGGCH ILSQVVRSSP GRASQR

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Mst1r Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its

specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Product Details Sterility: 0.22 µm filtered Endotoxin Level: Protein is endotoxin free. Grade: Crystallography grade **Target Details** MST1R Target: Alternative Name: Mst1r (MST1R Products) Background: Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to MST1 ligand. Regulates many physiological processes including cell survival, migration and differentiation. Ligand binding at the cell surface induces autophosphorylation of RON on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1 or the adapter GAB1. Recruitment of these downstream effectors by RON leads to the activation of several signaling cascades including the RAS-ERK, PI3 kinase-AKT, or PLCgamma-PKC. RON signaling activates the wound healing response by promoting epithelial cell migration, proliferation as well as survival at the wound site. Plays also a role in the innate immune response by regulating the migration and phagocytic activity of macrophages. Alternatively, RON can also promote signals such as cell migration and proliferation in response to growth factors other than MST1 ligand. Molecular Weight: 102.1 kDa Including tag. UniProt: Q62190 Pathways: **RTK Signaling Application Details** In addition to the applications listed above we expect the protein to work for functional studies **Application Notes:** as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though. Comment: Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn | International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com | Page 3/4 | Product datasheet for ABIN3135520 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

For Research Use only

Restrictions:

options with you in detail to assure that you receive your protein of interest.

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process