

Datasheet for ABIN3135579

TRIM24 Protein (AA 1-1051) (Strep Tag)



Go to Product page

_				
()	ve.	rv/	101	Λ

Quantity:	250 μg
Target:	TRIM24
Protein Characteristics:	AA 1-1051
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This TRIM24 protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), ELISA, Western Blotting (WB)

Product Details		
Brand:	AliCE®	
Sequence:	MEVAVEKAAA AAAPAGGPAA AAPSGENEAE SRQGPDSESG GEASRLNLLD TCAVCHQNIQ	
	SRVPKLLPCL HSFCQRCLPA PQRYLMLTAP ALGSAETPPP APAPAPAPGS PAGGPSPFAT	
	QVGVIRCPVC SQECAERHII DNFFVKDTTE VPSSTVEKSN QVCTSCEDNA EANGFCVECV	
	EWLCKTCIRA HQRVKFTKDH TVRQKEEVSP EAVGVTSQRP VFCPFHKKEQ LKLYCETCDK	
	LTCRDCQLLE HKEHRYQFIE EAFQNQKVII DTLITKLMEK TKYIKYTGNQ IQNRIIEINQ	
	NQKQVEQDIK VAIFTLMVEI NKKGKALLHQ LESLAKDHRM KLMQQQQEVA GLSKQLEHVM	
	HFSKWAVSSG SSTALLYSKR LITYRLRHLL RARCDASPVT NTTIQFHCDP SFWAQNIINL	
	GSLVIEDKES QPQMPKQNPV VEQSSQPPGG LPSNQLSKFP TQISLAQLRL QHIQQQVMAQ	
	RQQVQRRPAP VGLPNPRMQG PIQQPSISHQ HPPPRLINFQ NHSPKPNGPV LPPYPQQLRY	
	SPSQNVPRQT TIKPNPLQMA FLAQQAIKQW QISSVQAPPT TASSSSSTPS SPTITSAAGY	
	DGKAFSSPMI DLSAPVGGSY NLPSLPDIDC SSTIMLDNIA RKDTGVDHAQ PRPPSNRTVQ	

SPNSSVPSPG LAGPVTMTSV HPPIRSPSAS SVGSRGSSGS SSKPAGADST HKVPVVMLEP IRIKQENSGP PENYDFPVVI VKQESDEESR PQNTNYPRSI LTSLLLNSSQ SSASEETVLR SDAPDSTGDQ PGLHQENSSN GKSEWSDASQ KSPVHVGETR KEDDPNEDWC AVCQNGGELL CCEKCPKVFH LTCHVPTLTN FPSGEWICTF CRDLSKPEVD YDCDVPSHHS EKRKSEGLTK LTPIDKRKCE RLLLFLYCHE MSLAFQDPVP LTVPDYYKII KNPMDLSTIK KRLQEDYCMY TKPEDFVADF RLIFQNCAEF NEPDSEVANA GIKLESYFEE LLKNLYPEKR FPKVEFRHEA EDCKFSDDSD DDFVQPRKKR LKSTEDRQLL K

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- · State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

• The concentration of our recombinant proteins is measured using the absorbance at 280nm.

- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade:

custom-made

Target Details

Target:

TRIM24

Alternative Name:

Trim24 (TRIM24 Products)

Background:

Transcription intermediary factor 1-alpha (TIF1-alpha) (EC 2.3.2.27) (E3 ubiquitin-protein ligase Trim24) (RING-type E3 ubiquitin transferase TIF1-alpha) (Tripartite motif-containing protein 24), FUNCTION: Transcriptional coactivator that interacts with numerous nuclear receptors and coactivators and modulates the transcription of target genes. Interacts with chromatin depending on histone H3 modifications, having the highest affinity for histone H3 that is both unmodified at 'Lys-4' (H3K4me0) and acetylated at 'Lys-23' (H3K23ac) (By similarity). Has E3 protein-ubiquitin ligase activity. Promotes ubiquitination and proteasomal degradation of p53/TP53. Plays a role in the regulation of cell proliferation and apoptosis via its effects on p53/TP53 levels. Up-regulates ligand-dependent transcription activation by AR, GCR/NR3C1, thyroid hormone receptor (TR) and ESR1. Modulates transcription activation by retinoic acid (RA) receptors, such as RARA. Plays a role in regulating retinoic acid-dependent proliferation of hepatocytes. Required for normal transition from proliferating neonatal hepatocytes to quiescent adult hepatocytes. {ECO:0000250, ECO:0000269|PubMed:10610177, ECO:0000269|PubMed:16322096, ECO:0000269|PubMed:16880268,

ECO:0000269|PubMed:18026104, ECO:0000269|PubMed:19556538,

ECO:0000269|PubMed:19909775, ECO:0000269|PubMed:7744009}., FUNCTION:

Transcriptional coactivator that interacts with numerous nuclear receptors and coactivators and modulates the transcription of target genes. Interacts with chromatin depending on histone H3 modifications, having the highest affinity for histone H3 that is both unmodified at 'Lys-4' (H3K4me0) and acetylated at 'Lys-23' (H3K23ac). Has E3 protein-ubiquitin ligase activity. During the DNA damage response, participates in an autoregulatory feedback loop with TP53. Early in response to DNA damage, ATM kinase phosphorylates TRIM24 leading to its

ubiquitination and degradation. After sufficient DNA repair has occurred, TP53 activates TRIM24 transcription, ultimately leading to TRIM24-mediated TP53 ubiquitination and degradation (By similarity). Plays a role in the regulation of cell proliferation and apoptosis, at least in part via its effects on p53/TP53 levels. Up-regulates ligand-dependent transcription activation by AR, GCR/NR3C1, thyroid hormone receptor (TR) and ESR1. Modulates transcription activation by retinoic acid (RA) receptors, including RARA. Plays a role in regulating retinoic acid-dependent proliferation of hepatocytes (By similarity). Participates also in innate immunity by mediating the specific 'Lys-63'-linked ubiquitination of TRAF3 leading to activation of downstream signal transduction of the type I IFN pathway. Additionally, negatively regulates NLRP3/CASP1/IL-1beta-mediated pyroptosis and cell migration probably by ubiquitinating NLRP3 (By similarity). (ECO:0000250|UniProtKB:015164, ECO:0000269|PubMed:10610177, ECO:0000269|PubMed:16322096, ECO:0000269|PubMed:16880268, ECO:0000269|PubMed:18026104, ECO:0000269|PubMed:19556538, ECO:0000269|PubMed:19909775, ECO:0000269|PubMed:7744009}.

Molecular Weight:

116.7 kDa

UniProt:

Q64127

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment:

ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	12 months	