antibodies .- online.com





SLCO4C1 Protein (AA 1-722) (rho-1D4 tag)





Go to Product pag

Overview

Quantity:	1 mg
Target:	SLCO4C1 (OATP-H)
Protein Characteristics:	AA 1-722
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLCO4C1 protein is labelled with rho-1D4 tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

Product Details

Sequence:

MQGSKGIENP AFVPSSPGTP RRASASPSQV EVSAVASRNQ NGGSQPRESE EPQKSTEPSP PSSNPPASDE PPGSQLSELE EGPCGWRGFH PQCLQRCNTP QGFLLHYCLL ALTQGIVVNG LVNISISTIE KRYEMKSSLT GLISSSYDIS FCVLSLFVSF FGERGHKPRW LAFASFMIGL GALVFSLPHF FSGRYELGSI FEDTCLTRNS TRCSSSTSLL SNYFYVFVLG QLLLGTGGTP LYTLGTAFID DSVPTHKSSL YIGIGYSMSI LGPAIGYVLG GQLLTMYIDI AMGQSSDLTE DDPRWLGAWW IGFLLAWLFA WSLIMPFSCF PKHLPGTAKI QAGKTSQTHQ NNSTSFQHTD ENFGKSIKDF PTAVKNLMRN TVFICLVLST TSEALITTGF ATFLPKFIEN QFGLTSSFAA TLGGAVLIPG AALGQILGGV LVSKFKMKCK NTMKFALCTS GVALVLSFVF IYAKCENEPF AGVSESYNGT GEMGNLTAPC NANCNCLRSY YYPLCGSDGI QYFSPCFAGC LNSVSNRKPK VYYNCSCIER KITSTAESTD FEAKAGKCRT RCSNLPIFLG IFFITVIFTF MAGTPITVSI LRCVNHRHRS LALGVQFMLL RLLGTIPGPI IFGVIIDSTC VLWDVNECGI KGACWIYDNI KMAHMLVAIS VTCKVITIFF NGLAIVLYKP PPPGTEVSFQ SQNVIVSTIS VEEDLDKAEN EG

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Slco4c1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

- 1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Product Details Protein is endotoxin-free. Endotoxin Level: Grade: Crystallography grade Target Details SLCO4C1 (OATP-H) Target: Slco4c1 (OATP-H Products) Alternative Name: Background: Organic anion transporter, capable of transporting pharmacological substances such as digoxin, ouabain, thyroxine, methotrexate and cAMP. May participate in the regulation of membrane transport of ouabain. Involved in the uptake of the dipeptidyl peptidase-4 inhibitor sitagliptin and hence may play a role in its transport into and out of renal proximal tubule cells. May be involved in the first step of the transport pathway of digoxin and various compounds into the urine in the kidney. May be involved in sperm maturation by enabling directed movement of organic anions and compounds within or between cells. This ion-transporting process is important to maintain the strict epididymal homeostasis necessary for sperm maturation. May have a role in secretory functions since seminal vesicle epithelial cells are assumed to secrete proteins involved in decapacitation by modifying surface proteins to facilitate the acquisition of the ability to fertilize the egg. {ECO:0000250|UniProtKB:Q6ZQN7, ECO:0000269|PubMed:16641146}. Molecular Weight: 79.6 kDa Including tag. UniProt: Q8BGD4 **Application Details Application Notes:** In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though. Comment: Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible

For Research Use only

Restrictions:

options with you in detail to assure that you receive your protein of interest.

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process