antibodies .- online.com





HIF1AN Protein (AA 2-349) (His tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	HIF1AN
Protein Characteristics:	AA 2-349
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This HIF1AN protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA, Crystallization (Crys)

Product Details

Sequence:

AATAAEVAAS GSGEAREEAE APGPAWDESQ LRSYSFPTRP IPRLSQSDPR AEELIENEEP VVLTDTNLVY PALKWDLEYL QENIGNGDFS VYSASTHKFL YYDEKKMGNF QNFKPRSNRE EIKFHEFVEK LQAIQQRGGE ERLYLQQTLN DTVGRKIVMD FLGFNWNWIN KQQGKRGWGQ LTSNLLLIGM EGNVTPAHYD EQQNFFAQIK GHKRCILFPP DQFECLYPYP VHHPCDRQSQ VDFDNPDYER FPNFRNVVGY ETVVGPGDVL YIPMYWWHHI ESLLNGGITI TVNFWYKGAP TPKRIEYPLK AHQKVAIMRN IEKMLGEALG NPQEVGPLLN TMIKGRYN

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Hif1an Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

Target Details

Target:	HIF1AN
Alternative Name:	Hif1an (HIF1AN Products)
Background:	Hydroxylates HIF-1 alpha at 'Asp-799' in the C-terminal transactivation domain (CAD).

. d. got 2 otalio	
	Functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents
	interaction of HIF-1 with transcriptional coactivators including Cbp/p300-interacting
	transactivator. Involved in transcriptional repression through interaction with HIF1A, VHL and
	histone deacetylases. Hydroxylates specific Asn residues within ankyrin repeat domains (ARD)
	of NFKB1, NFKBIA, NOTCH1, ASB4, PPP1R12A and several other ARD-containing proteins. Also
	hydroxylates Asp and His residues within ARDs of ANK1 and TNKS2, respectively. Negatively
	regulates NOTCH1 activity, accelerating myogenic differentiation (By similarity). Positively
	regulates ASB4 activity, promoting vascular differentiation. {ECO:0000250 UniProtKB:Q9NWT6,
	ECO:0000269 PubMed:17636018}.
Molecular Weight:	41.1 kDa Including tag.
UniProt:	Q8BLR9
Pathways:	Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the
	recombinant protein with the default tag will be insoluble our protein lab may suggest a higher
	molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible
	options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.

-80 °C

Store at -80°C.

Unlimited (if stored properly)

Storage:

Expiry Date:

Storage Comment:




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process