

Datasheet for ABIN3136553  
**SIRT5 Protein (AA 37-310) (His tag)**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	1 mg
Target:	SIRT5
Protein Characteristics:	AA 37-310
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SIRT5 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

## Product Details

Sequence:	<p>SSNMADFRKC FANAKHIAII SGAGVSAESG VPTFRGAGGY WRKWQAQDLA TPQAFARNPS QVWEFYHYRR EVMRSKEPNP GHIAIAQCEA RLRDQGRRVV VITQNIDELH RKAGTKNLLE IHGTLFKTRC TSCGTVAENY RSPICPALAG KGAPEPETQD ARIPVDKLPR CEEAGCGLL RPHVVWFGEN LDPAILEEVD RELALCDLCL VVGTSVVYP AAMFAPQVAS RGVPAEFNM ETTPATDRFR FHFPGPCGKT LPEALAPHET ERTS</p> <p><b>Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.</b></p>
Characteristics:	<ul style="list-style-type: none"><li>• Made in Germany - from design to production - by highly experienced protein experts.</li><li>• Mouse Sirt5 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.</li><li>• State-of-the-art algorithm used for plasmid design (Gene synthesis).</li></ul> <p>This protein is a made to order protein and will be made for the first time for your order. Our</p>

## Product Details

---

experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the ExPASy's protparam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in baculovirus infected SF9 insect cells:  1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.  2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Protein is endotoxin free.
Grade:	Crystallography grade

## Target Details

---

Target:	SIRT5
Alternative Name:	Sirt5 ( <a href="#">SIRT5 Products</a> )
Background:	NAD-dependent lysine demalonylase, desuccinylase and deglutarylase that specifically removes malonyl, succinyl and glutaryl groups on target proteins (PubMed:23806337,

## Target Details

PubMed:21908771, PubMed:22076378, PubMed:24315375, PubMed:24703693). Activates CPS1 and contributes to the regulation of blood ammonia levels during prolonged fasting: acts by mediating desuccinylation and deglutarylation of CPS1, thereby increasing CPS1 activity in response to elevated NAD levels during fasting (PubMed:19410549, PubMed:24703693). Activates SOD1 by mediating its desuccinylation, leading to reduced reactive oxygen species. Modulates ketogenesis through the desuccinylation and activation of HMGCS2 (PubMed:24315375). Has weak NAD-dependent protein deacetylase activity, however this activity may not be physiologically relevant in vivo. Can deacetylate cytochrome c (CYCS) and a number of other proteins in vitro such as Uox (PubMed:23085393). {ECO:0000269|PubMed:19410549, ECO:0000269|PubMed:21908771, ECO:0000269|PubMed:22076378, ECO:0000269|PubMed:23085393, ECO:0000269|PubMed:23806337, ECO:0000269|PubMed:24315375, ECO:0000269|PubMed:24703693}.

Molecular Weight: 31.1 kDa Including tag.

UniProt: [Q8K2C6](#)

Pathways: [SARS-CoV-2 Protein Interactome](#)

## Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: 100 mM NaCl, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

## Handling

---

Storage Comment: Store at -80°C.

Expiry Date: Unlimited (if stored properly)

## Images

---



**Image 1.** „Crystallography Grade“ protein due to multi-step, protein-specific purification process