antibodies .- online.com





PAF1/PD2 Protein (AA 1-535) (Strep Tag)



Overview

Quantity:	1 mg
Target:	PAF1/PD2 (PAF1)
Protein Characteristics:	AA 1-535
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This PAF1/PD2 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Sequence:

MAPTIQTQAQ REDGHRPNSH RTLPERSGVV CRVKYCNSLP DIPFDPKFIT YPFDQNRFVQ
YKATSLEKQH KHDLLTEPDL GVTIDLINPD TYRIDPNVLL DPADEKLLEE EIQAPTSSKR
SQQHAKVVPW MRKTEYISTE FNRYGISNEK PEVKIGVSVK QQFTEEEIYK DRDSQITAIE
KTFEDAQKSI SQHYSKPRVT PVEVMPVFPD FKMWINPCAQ VIFDSDPAPK DTSGAAALEM
MSQAMIRGMM DEEGNQFVAY FLPVEETLKK RKRDQEEEMD YAPDDVYDYK IAREYNWNVK
NKASKGYEEN YFFIFREGDG VYYNELETRV RLSKRRAKAG VQSGTNALLV VKHRDMNEKE
LEAQEARKAQ LENHEPEEEE EEEMEAEEKE AGGSDEEQEK GSSSEKEGSE DEHSGSESDR
EEGDRDEASD KSGSGEDESS EDEARAARDK EEIFGSDADS EDDADSDDED RGQAHRGSDN
DSDSGSDGGG QRSRSQSRSR SRSASPFPSG SEHSAQEDGS EAAASDSSEA DSDSD

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and

Product Details

Product Details	
	Western blot.
Purity:	≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Target Details	
Target:	PAF1/PD2 (PAF1)
Alternative Name:	Paf1 (PAF1 Products)
Background:	RNA polymerase II-associated factor 1 homolog, FUNCTION: Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1), UB2A/B-mediated H2E ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. Connects PAF1C with the RNF20/40 E3 ubiquitin-protein ligase complex. Involved in polyadenylation of mRNA
Molecular Weight:	precursors (By similarity). {ECO:0000250, ECO:0000269 PubMed:19345177}. 60.5 kDa
UniProt:	Q8K2T8
Pathways:	Cellular Response to Molecule of Bacterial Origin, Stem Cell Maintenance
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from

Application Details

Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)