

Datasheet for ABIN3136787

DDX1 Protein (AA 1-740) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	DDX1
Protein Characteristics:	AA 1-740
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This DDX1 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Brand:	AliCE®
Sequence:	MAAFSEMGVM PEIAQAVEEM DWLLPTDIQA ESIPLILGGG DVLMAAETGS GKTGAFSIPV IQIVYETLKD QQEGKKGKTT IKTGASVLNK WQMNPYDRGS AFAIGSDGLC CQSREVKWEH GCRGTRGLLK GKHYYEVSCH DQGLCRVGWS TMQASLDLGT DKFGFGFGGT GK KSHNKQFD NYGEEFTMHD TIGCYLDIDK GHVKFSKNGK DLGLAFEIPA HIKNQALFPA CVLKNAELKF NFGEEEFKFP PKDGFVALSK APDNYIVKSQ HTGNAQVSQT KFLPNAPKAL IVEPSRELAE QTLNNVKQFK KYIDNPKLRE LLIIGGVAAR DQLSVLDNGV DIVVGTPGRL DDLVSTGKLN LSQVRFLVLD EADGLLSQGY SDFINRMHNQ IPQITCDGKR LQVIVCSATL HSFDVKKLSE KIMHFPTWVD LKGEDSVPDT VHHVVVPVNP KTDKLWERLG KNHIRTDDVH AKDNTRPGAN SPEMWSEAIK ILKGEYAVRA IKEHKMDQAI IFCRTKIDCD NLEQYFMQQG GGPDKKGHQF SCVCLHGDRK PHERKQNLER FKKGDVRFLLI CTDVAARGID IHGVPYVINV TLPDEKQNYV HRIGRVGRAE RMGLAISLVA TEKEKVWYHV CSNRGKGCYN TRLKEDGGCT IWYNEMQLLS

EIEEHLNCTI SQVEPDIKVP VDEFDGKVTY GQKRAAGGGN YKGHVDVLAP TVQELAALEK
EAQTSFLHLG YLPNQLFRTF

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Product Details

Purity: > 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade: custom-made

Target Details

Target: DDX1

Alternative Name: Ddx1 ([DDX1 Products](#))

Background: ATP-dependent RNA helicase DDX1 (EC 3.6.4.13) (DEAD box protein 1),FUNCTION: Acts as an ATP-dependent RNA helicase, able to unwind both RNA-RNA and RNA-DNA duplexes. Possesses 5' single-stranded RNA overhang nuclease activity. Possesses ATPase activity on various RNA, but not DNA polynucleotides. May play a role in RNA clearance at DNA double-strand breaks (DSBs), thereby facilitating the template-guided repair of transcriptionally active regions of the genome. Together with RELA, acts as a coactivator to enhance NF-kappa-B-mediated transcriptional activation (By similarity). Acts as a positive transcriptional regulator of cyclin CCND2 expression (PubMed:19398953). Binds to the cyclin CCND2 promoter region (PubMed:19398953). Associates with chromatin at the NF-kappa-B promoter region via association with RELA. Binds to poly(A) RNA. May be involved in 3'-end cleavage and polyadenylation of pre-mRNAs. Component of the tRNA-splicing ligase complex required to facilitate the enzymatic turnover of catalytic subunit RTCB: together with archease (ZBTB80S), acts by facilitating the guanylylation of RTCB, a key intermediate step in tRNA ligation (By similarity). Component of a multi-helicase-TICAM1 complex that acts as a cytoplasmic sensor of viral double-stranded RNA (dsRNA) and plays a role in the activation of a cascade of antiviral responses including the induction of pro-inflammatory cytokines via the adapter molecule TICAM1 (PubMed:21703541). Specifically binds (via helicase ATP-binding domain) on both short and long poly(I:C) dsRNA (PubMed:21703541). {ECO:0000250|UniProtKB:Q92499, ECO:0000269|PubMed:19398953, ECO:0000269|PubMed:21703541}.

Molecular Weight: 82.5 kDa

UniProt: [Q91VR5](#)

Pathways: [Ribonucleoprotein Complex Subunit Organization](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Application Details

Comment:	<p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!</p>
----------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Buffer:	<p>The buffer composition is at the discretion of the manufacturer.</p> <p>Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.</p>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months