

Datasheet for ABIN3136803
PRKAG2 Protein (AA 1-566) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	PRKAG2
Protein Characteristics:	AA 1-566
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This PRKAG2 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Sequence: MGSAAMDTKK KKEVSSPGGS SGKKNPSLKR RSLRVHIPDL SSFAMPLLDG DVENSEKHSS
RKVDSPFSSG SPSRGLFSRG PQPRPSSPVS APVRPKTSPG SPKTVFPFSY QESPPRSPRR
MSFSGIFRSS SKESPNSNP STSPGGIRFF SRSRKTSSVS SSPSTPTQVT KQHPPLESY
KQEPERPESE IYASSSPPDT GQRFCLAFQS PARPPLASPT YHAPLRTAVL AAAPGPAEAG
MLEKLEFQEE EDSSEGVYMR FMRSHKCYDI VPTSSKLVVF DDTLQVKKAF FALVANGVRA
APLWESKKQS FVGMLTITDF INILHRYYS PMVQIYELEE HKIETWRELY LQETFKPLVN
ISPDASLFDA VYSLIKNKIH RLPVIDPISG NALYILTHKR ILKFLQLFMS DMPKPAPFMKQ
NLDELGIGTY HNIAFIHPDT PIIKALNIFV ERRISALPVV DESGKVVDIY SKFDVINLAA EKTYNNLDIT
VTQALQHRSQ YFEGVVKCSK LETLETIVDR IVRAEVHRLV VVNEADSVG IISLSDILQA
LILTPAGAKQ KETETE

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Target Details

Target:	PRKAG2
Alternative Name:	Prkag2 (PRKAG2 Products)
Background:	5'-AMP-activated protein kinase subunit gamma-2 (AMPK gamma2) (AMPK subunit gamma-2),FUNCTION: AMP/ATP-binding subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton, probably by indirectly activating myosin. Gamma non-catalytic subunit mediates binding to AMP, ADP and ATP, leading to activate or inhibit AMPK: AMP-binding results in allosteric activation of alpha catalytic subunit (PRKAA1 or PRKAA2) both by inducing phosphorylation and preventing dephosphorylation of catalytic subunits. ADP also stimulates phosphorylation, without stimulating already phosphorylated catalytic subunit. ATP promotes dephosphorylation of catalytic subunit, rendering the AMPK enzyme inactive. {ECO:0000250 UniProtKB:Q9UGJ0}.
Molecular Weight:	62.9 kDa
UniProt:	Q91WG5
Pathways:	AMPK Signaling , Cellular Glucan Metabolic Process , Ribonucleoside Biosynthetic Process , Regulation of Carbohydrate Metabolic Process , Warburg Effect

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce

Application Details

something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.
Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months
