

# Datasheet for ABIN3136834 Arrestin 3 Protein (AA 1-410) (Strep Tag)



Overview

Quantity:	1 mg
Target:	Arrestin 3 (ARRB2)
Protein Characteristics:	AA 1-410
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This Arrestin 3 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

## Product Details

Brand:	AliCE®
Sequence:	MGEKPGTRVF KKSSPNCKLT VYLGKRDFVD HLDKVDPVDG VVLVDPDYLK DRKVFVTLTC
	AFRYGREDLD VLGLSFRKDL FIATYQAFPP MPNPPRPPTR LQDRLLKKLG QHAHPFFFTI
	PQNLPCSVTL QPGPEDTGKA CGVDFEIRAF CAKSIEEKSH KRNSVRLIIR KVQFAPETPG
	PQPSAETTRH FLMSDRRSLH LEASLDKELY YHGEPLNVNV HVTNNSAKTV KKIRVSVRQY
	ADICLFSTAQ YKCPVAQLEQ DDQVSPSSTF CKVYTITPLL SDNREKRGLA LDGQLKHEDT
	NLASSTIVKE GANKEVLGIL VSYRVKVKLV VSRGGDVSVE LPFVLMHPKP HDHITLPRPQ
	SAPRETDVPV DTNLIEFDTN YATDDDIVFE DFARLRLKGM KDDDCDDQFC
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/5 | Product datasheet for ABIN3136834 | 02/25/2025 | Copyright antibodies-online. All rights reserved.

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made
Target Details	
Target:	Arrestin 3 (ARRB2)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/5 | Product datasheet for ABIN3136834 | 02/25/2025 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	Arrb2 (ARRB2 Products)
Background:	Beta-arrestin-2 (Arrestin beta-2),FUNCTION: Functions in regulating agonist-mediated G-protein
	coupled receptor (GPCR) signaling by mediating both receptor desensitization and
	resensitization processes. During homologous desensitization, beta-arrestins bind to the GPRK-
	phosphorylated receptor and sterically preclude its coupling to the cognate G-protein, the
	binding appears to require additional receptor determinants exposed only in the active receptor
	conformation. The beta-arrestins target many receptors for internalization by acting as
	endocytic adapters (CLASPs, clathrin-associated sorting proteins) and recruiting the GPRCs to
	the adapter protein 2 complex 2 (AP-2) in clathrin-coated pits (CCPs). However, the extent of
	beta-arrestin involvement appears to vary significantly depending on the receptor, agonist and
	cell type. Internalized arrestin-receptor complexes traffic to intracellular endosomes, where they
	remain uncoupled from G-proteins. Two different modes of arrestin-mediated internalization
	occur. Class A receptors, like ADRB2, OPRM1, ENDRA, D1AR and ADRA1B dissociate from beta-
	arrestin at or near the plasma membrane and undergo rapid recycling. Class B receptors, like
	AVPR2, AGTR1, NTSR1, TRHR and TACR1 internalize as a complex with arrestin and traffic with
	it to endosomal vesicles, presumably as desensitized receptors, for extended periods of time.
	Receptor resensitization then requires that receptor-bound arrestin is removed so that the
	receptor can be dephosphorylated and returned to the plasma membrane. Mediates
	endocytosis of CCR7 following ligation of CCL19 but not CCL21. Involved in internalization of
	P2RY1, P2RY4, P2RY6 and P2RY11 and ATP-stimulated internalization of P2RY2. Involved in
	phosphorylation-dependent internalization of OPRD1 and subsequent recycling or degradation.
	Involved in ubiquitination of IGF1R. Beta-arrestins function as multivalent adapter proteins that
	can switch the GPCR from a G-protein signaling mode that transmits short-lived signals from
	the plasma membrane via small molecule second messengers and ion channels to a beta-
	arrestin signaling mode that transmits a distinct set of signals that are initiated as the receptor
	internalizes and transits the intracellular compartment. Acts as a signaling scaffold for MAPK
	pathways such as MAPK1/3 (ERK1/2) and MAPK10 (JNK3). ERK1/2 and JNK3 activated by the
	beta-arrestin scaffold are largely excluded from the nucleus and confined to cytoplasmic
	locations such as endocytic vesicles, also called beta-arrestin signalosomes. Acts as a
	signaling scaffold for the AKT1 pathway. GPCRs for which the beta-arrestin-mediated signaling
	relies on both ARRB1 and ARRB2 (codependent regulation) include ADRB2, F2RL1 and PTH1R.
	For some GPCRs the beta-arrestin-mediated signaling relies on either ARRB1 or ARRB2 and is
	inhibited by the other respective beta-arrestin form (reciprocal regulation). Increases ERK1/2
	signaling in AGTR1- and AVPR2-mediated activation (reciprocal regulation). Involved in CCR7-
	mediated ERK1/2 signaling involving ligand CCL19. Is involved in type-1A angiotensin II

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/5 | Product datasheet for ABIN3136834 | 02/25/2025 | Copyright antibodies-online. All rights reserved. receptor/AGTR1-mediated ERK activity. Is involved in type-1A angiotensin II receptor/AGTR1mediated MAPK10 activity. Is involved in dopamine-stimulated AKT1 activity in the striatum by disrupting the association of AKT1 with its negative regulator PP2A. Involved in AGTR1mediated chemotaxis. Appears to function as signaling scaffold involved in regulation of MIP-1beta-stimulated CCR5-dependent chemotaxis. Involved in attenuation of NF-kappa-B-dependent transcription in response to GPCR or cytokine stimulation by interacting with and stabilizing CHUK. Suppresses UV-induced NF-kappa-B-dependent activation by interacting with CHUK. The function is promoted by stimulation of ADRB2 and dephosphorylation of ARRB2. Involved in IL8-mediated granule release in neutrophils (By similarity). Involved in p53/TP53-mediated apoptosis by regulating MDM2 and reducing the MDM2-mediated degradation of p53/TP53. May serve as nuclear messenger for GPCRs. Upon stimulation of OR1D2, may be involved in regulation of gene expression during the early processes of fertilization. Also involved in regulation of receptors other than GPCRs. Involved in endocytosis of TGFBR2 and TGFBR3 and down-regulates TGF-beta signaling such as NF-kappa-B activation. Involved in endocytosis of low-density lipoprotein receptor/LDLR. Involved in endocytosis of smoothened homolog/Smo, which also requires GRK2. Involved in endocytosis of SLC9A5. Involved in endocytosis of ENG and subsequent TGF-beta-mediated ERK activation and migration of epithelial cells. Involved in Toll-like receptor and IL-1 receptor signaling through the interaction with TRAF6 which prevents TRAF6 autoubiquitination and oligomerization required for activation of NF-kappa-B and JUN. Involved in insulin resistance by acting as insulin-induced signaling scaffold for SRC, AKT1 and INSR. Involved in regulation of inhibitory signaling of natural killer cells by recruiting PTPN6 and PTPN11 to KIR2DL1. Involved in the internalization of the atypical chemokine receptor ACKR3 (By similarity). Acts as an adapter protein coupling FFAR4 receptor to specific downstream signaling pathways, as well as mediating receptor endocytosis. During the activation step of NLRP3 inflammasome, directly associates with NLRP3 leading to inhibition of pro-inflammatory cytokine release and inhibition of inflammation. {ECO:0000250, ECO:0000250|UniProtKB:P32121, ECO:0000269|PubMed:12944399, ECO:0000269|PubMed:16051150, ECO:0000269|PubMed:17540773, ECO:0000269|PubMed:18337459, ECO:0000269|PubMed:18604210, ECO:0000269|PubMed:19122674}.

Molecular Weight:	46.3 kDa
UniProt:	Q91YI4
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway, Regulation of Intracellular Steroid Hormone Receptor Signaling, cAMP Metabolic Process, Myometrial Relaxation and Contraction, Regulation of Leukocyte Mediated Immunity, Synaptic Membrane, Regulation of G-Protein

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/5 | Product datasheet for ABIN3136834 | 02/25/2025 | Copyright antibodies-online. All rights reserved.

## Target Details

Coupled Receptor Protein Signaling, CXCR4-mediated Signaling Events, Phototransduction, Thromboxane A2 Receptor Signaling

### **Application Details**

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol <b>Might differ depending on protein.</b>
Handling Advice:	Avoid repeated freeze-thaw cycles.

Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 5/5 | Product datasheet for ABIN3136834 | 02/25/2025 | Copyright antibodies-online. All rights reserved.