

Datasheet for ABIN3136924 CDK9 Protein (AA 1-372) (Strep Tag)



Overview

Quantity:	250 µg
Target:	CDK9
Protein Characteristics:	AA 1-372
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CDK9 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), ELISA, SDS-PAGE (SDS)

Product Details

Brand:	AliCE®
Sequence:	MAKQYDSVEC PFCDEVTKYE KLAKIGQGTF GEVFKAKHRQ TGQKVALKKV LMENEKEGFP
	ITALREIKIL QLLKHENVVN LIEICRTKAS PYNRCKGSIY LVFDFCEHDL AGLLSNVLVK
	FTLSEIKRVM QMLLNGLYYI HRNKILHRDM KAANVLITRD GVLKLADFGL ARAFSLAKNS
	QPNRYTNRVV TLWYRPPELL LGERDYGPPI DLWGAGCIMA EMWTRSPIMQ GNTEQHQLAL
	ISQLCGSITP EVWPNVDKYE LFEKLELVKG QKRKVKDRLK AYVRDPYALD LIDKLLVLDP
	AQRIDSDDAL NHDFFWSDPM PSDLKGMLST HLTSMFEYLA PPRRKGSQIT QQSTNQSRNP
	ATTNQTEFER VF
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN3136924 | 02/26/2025 | Copyright antibodies-online. All rights reserved.

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
custom-made
CDK9

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3136924 | 02/26/2025 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	Cdk9 (CDK9 Products)
Background:	Cyclin-dependent kinase 9 (EC 2.7.11.22) (EC 2.7.11.23) (Cell division protein kinase
	9),FUNCTION: Protein kinase involved in the regulation of transcription. Member of the cyclin-
	dependent kinase pair (CDK9/cyclin-T) complex, also called positive transcription elongation
	factor b (P-TEFb), which facilitates the transition from abortive to productive elongation by
	phosphorylating the CTD (C-terminal domain) of the large subunit of RNA polymerase II (RNAP
	II) POLR2A, SUPT5H and RDBP. This complex is inactive when in the 7SK snRNP complex form.
	Phosphorylates EP300, MYOD1, RPB1/POLR2A and AR and the negative elongation factors
	DSIF and NELFE. Regulates cytokine inducible transcription networks by facilitating promoter
	recognition of target transcription factors (e.g. TNF-inducible RELA/p65 activation and IL-6-
	inducible STAT3 signaling). Promotes RNA synthesis in genetic programs for cell growth,
	differentiation and viral pathogenesis. P-TEFb is also involved in cotranscriptional histone
	modification, mRNA processing and mRNA export. Modulates a complex network of chromatin
	modifications including histone H2B monoubiquitination (H2Bub1), H3 lysine 4 trimethylation
	(H3K4me3) and H3K36me3, integrates phosphorylation during transcription with chromatin
	modifications to control co-transcriptional histone mRNA processing. The CDK9/cyclin-K
	complex has also a kinase activity towards CTD of RNAP II and can substitute for CDK9/cyclin-
	T P-TEFb in vitro. Replication stress response protein, the CDK9/cyclin-K complex is required
	for genome integrity maintenance, by promoting cell cycle recovery from replication arrest and
	limiting single-stranded DNA amount in response to replication stress, thus reducing the
	breakdown of stalled replication forks and avoiding DNA damage. In addition, probable function
	in DNA repair of isoform 2 via interaction with KU70/XRCC6. Promotes cardiac myocyte
	enlargement. RPB1/POLR2A phosphorylation on 'Ser-2' in CTD activates transcription. AR
	phosphorylation modulates AR transcription factor promoter selectivity and cell growth. DSIF
	and NELF phosphorylation promotes transcription by inhibiting their negative effect. The
	phosphorylation of MYOD1 enhances its transcriptional activity and thus promotes muscle
	differentiation. Catalyzes phosphorylation of KAT5, promoting KAT5 recruitment to chromatin
	and histone acetyltransferase activity. {ECO:0000250 UniProtKB:P50750}.
Molecular Weight:	42.8 kDa
UniProt:	Q99J95
Pathways:	Cell Division Cycle

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN3136924 | 02/26/2025 | Copyright antibodies-online. All rights reserved.

Application Details	
	as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	 ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months