antibodies .- online.com





PPID Protein (AA 2-370) (His tag)



Go to Product page

Overview

Quantity:	1 mg	
Target:	PPID	
Protein Characteristics:	AA 2-370	
Origin:	Mouse	
Source:	Insect Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This PPID protein is labelled with His tag.	
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA, Crystallization (Crys)	

Product Details

Sequence:

SHASPAAKPS NSKNPRVFFD VDIGGERVGR IVLELFADIV PKTAENFRAL CTGEKGTGST
TGKPLHFKGC PFHRIIKKFM IQGGDFSNQN GTGGESIYGE KFEDENFHYK HDREGLLSMA
NAGPNTNGSQ FFITTVPTPH LDGKHVVFGQ VIKGLGVART LENVEVNGEK PAKLCVIAEC
GELKEGDDWG IFPKDGSGDS HPDFPEDADI DLKDVDKILL ISEDLKNIGN TFFKSQNWEM
AIKKYAKVLR YVDSSKAVIE KADRSRLQPI ALSCVLNIGA CKLKMSNWQG AIDSCLEALE
MDPSNTKALY RKAQGWQGLK EYDQALADLK KAQEIAPGDK AIQAELLKVK QMIKAQKDKE
KAVYAKMFA

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Ppid Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility: 0.22 µm filtered

Endotoxin Level: Protein is endotoxin free.

Grade: Crystallography grade

Target Details

Target: PPID

Alternative Name: Ppid (PPID Products)

Target Details

PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline	
imidic peptide bonds in oligopeptides. Proposed to act as a co-chaperone in HSP90 complexes	
such as in unligated steroid receptors heterocomplexes. Different co-chaperones seem to	
compete for association with HSP90 thus establishing distinct HSP90-co-chaperone-receptor	
complexes with the potential to exert tissue-specific receptor activity control. May have a	
preference for estrogen receptor complexes and is not found in glucocorticoid receptor	
complexes. May be involved in cytoplasmic dynein-dependent movement of the receptor from	
the cytoplasm to the nucleus. May regulate MYB by inhibiting its DNA-binding activity. Involved	
in regulation of AHR signaling by promoting the formation of the AHR:ARNT dimer, the function	
is independent of HSP90 but requires the chaperone activity region. Involved in regulation of UN	
radiation-induced apoptosis. {ECO:0000269 PubMed:18771283}.	
41.6 kDa Including tag.	
Q9CR16	
Nuclear Hormone Receptor Binding	
In addition to the applications listed above we expect the protein to work for functional studies	
as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee	
though.	
Protein has not been tested for activity yet. In cases in which it is highly likely that the	
recombinant protein with the default tag will be insoluble our protein lab may suggest a higher	
molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible	
options with you in detail to assure that you receive your protein of interest.	
For Research Use only	
Liquid	
100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Avoid repeated freeze-thaw cycles.	
-80 °C	
Store at -80°C.	

1	1 11	١٠
\vdash	land	unc
	ıarıa	11110

Expiry Date:

Unlimited (if stored properly)