# antibodies .- online.com





# SNW1 Protein (AA 2-536) (His tag)



## Overview

Quantity:	1 mg
Target:	SNW1
Protein Characteristics:	AA 2-536
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SNW1 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

## **Product Details**

Sequence:

ALTSFLPAPT QLSQDQLEAE ERARSQRSLQ TSLVSSRREP PPYGYRKGWI PRLLEDFGDG GAFPEIHVAQ YPLDMGRKKK MSNALAIQVD PEGKIKYDAI ARQGQSKDKV IYSKYTDLVP KEVMNADDPD LQRPDEEAIK EITEKTRVAL EKSVSQKVAA AMPVRAADKL APAQYIRYTP SQQGVAFNSG AKQRVIRMVE MQKEPMEPPR FKINKKIPRG PPSPPAPVMH SPSRKMTVKE QQEWKIPPCI SNWKNAKGYT IPIDKRLAAD GRGLQTVHIN ENFAKLAEAL YIADRKAREA VEMRAQVERK MAQKEKEKHE EKLREMAQKA RERRAGIKTH VEKEDGEARE RDEIRHDRRK ERQHDRNLSR AAPDKRSKLQ RNENRDISEV IALGVPNPRT SNEVQYDQRL FNQSKGMDSG FAGGEDEIYN VYDQAWRGGK DMAQSIYRPS KNLDKDMYGD DLEARIKTNR FVPDKEFSGS DRKQRGREGP VQFEEDPFGL DKFLEEAKQH GGSKRPSDSS RPKEHEHEGK KRRKE Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

#### Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Snw1 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

### Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Endotoxin has not been removed. Please contact us if you require endotoxin removal.
Grade:	Crystallography grade

# **Target Details**

Target:	SNW1
Alternative Name:	Snw1 (SNW1 Products)
Background:	Involved in transcriptional regulation. Modulates TGF-beta-mediated transcription via
	association with SMAD proteins, MYOD1-mediated transcription via association with PABPN1,
	RB1-mediated transcriptional repression, and retinoid-X receptor (RXR)- and vitamin D receptor
	(VDR)-dependent gene transcription in a cell line-specific manner probably involving
	coactivators NCOA1 and GRIP1. Is involved in NOTCH1-mediated transcriptional activation.
	Binds to multimerized forms of Notch intracellular domain (NICD) and is proposed to recruit
	transcriptional coactivators such as MAML1 to form an intermediate preactivation complex
	which associates with DNA-bound CBF-1/RBPJ to form a transcriptional activation complex by
	releasing SNW1 and redundant NOTCH1 NICD. Functions as a splicing factor in pre-mRNA
	splicing. Is required in the specific splicing of CDKN1A pre-mRNA, the function probbaly
	involves the recruitment of U2AF2 to the mRNA. Is proposed to recruit PPIL1 to the
	spliceosome. May be involved in cyclin-D1/CCND1 mRNA stability through the SNARP complex
	which associates with both the 3'end of the CCND1 gene and its mRNA (By similarity).
	{ECO:0000250}.
Molecular Weight:	62.3 kDa Including tag.
UniProt:	Q9CSN1
Pathways:	Retinoic Acid Receptor Signaling Pathway
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee
	though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the
	recombinant protein with the default tag will be insoluble our protein lab may suggest a higher
	molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible
	options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
	Liquid

# Handling

Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)