# antibodies .- online.com





Datasheet for ABIN3137087

# RHOH Protein (AA 1-188) (His tag)



**Image** 



#### Overview

Quantity:	1 mg
Target:	RHOH
Protein Characteristics:	AA 1-188
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This RHOH protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)
Product Details	

	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	KLFSINEC
	REVGPHRASC INAIEGKRLA QDVRAKGYLE CSALSNRGVQ QVFECAVRTA VNQARRRNRR
	GNDAFRSIRP LSYQQADVVL MCYSVANHNS FLNLKNKWIS EIRSNLPCTP VLVVATQTDQ
Sequence:	MLSSIKCVLV GDSAVGKTSL LVRFTSETFP EAYKPTVYEN TGVDVFMDGI QISLGLWDTA

special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Rhoh Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

## **Target Details**

Target:	RHOH
Alternative Name:	Rhoh (RHOH Products)
Background:	Binds GTP but lacks intrinsic GTPase activity and is resistant to Rho-specific GTPase-activating
	proteins. Inhibits the activation of NF-kappa-B by TNF and IKKB and the activation of CRK/p38
	by TNF. Inhibits activities of RAC1, RHOA and CDC42. Negatively regulates leukotriene

production in neutrophils (By similarity). Negative regulator of hematopoietic progenitor cell proliferation, survival and migration. Critical regulator of thymocyte development and T-cell antigen receptor (TCR) signaling by mediating recruitment and activation of ZAP70. Required for phosphorylation of CD3Z, membrane translocation of ZAP70 and subsequent activation of the ZAP70-mediated pathways. Essential for efficient beta-selection and positive selection by promoting the ZAP70-dependent phosphorylation of the LAT signalosome during pre-TCR and TCR signaling. Crucial for thymocyte maturation during DN3 to DN4 transition and during positive selection. Plays critical roles in mast cell function by facilitating phosphorylation of SYK in Fc epsilon RI-mediated signal transduction. Essential for the phosphorylation of LAT, LCP2, PLCG1 and PLCG2 and for Ca(2+) mobilization in mast cells. {ECO:0000250, ECO:0000269|PubMed:15494435, ECO:0000269|PubMed:17028588, ECO:0000269|PubMed:17119112, ECO:0000269|PubMed:19124738}.

Molecular Weight:

21.9 kDa Including tag.

UniProt:

Q9D3G9

# **Application Details**

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.

Comment:

Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process