antibodies .- online.com





SNX5 Protein (AA 2-404) (His tag)







Overview

Quantity:	1 mg
Target:	SNX5
Protein Characteristics:	AA 2-404
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SNX5 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)

Product Details

Sequence:

AAVPELLEQQ EEDRSKLRSV SVDLNVDPSL QIDIPDALSE RDKVKFTVHT KTTLSTFQSP EFSVTRQHED FVWLHDTLTE TTDYAGLIIP PAPTKPDFDG PREKMQKLGE GEGSMTKEEF AKMKQELEAE YLAVFKKTVS THEVFLQRLS SHPVLSKDRN FHVFLEYDQD LSVRRKNTKE MFGGFFKSVV KSADEVLFSG VKEVDDFFEQ EKNFLINYYN RIKDSCAKAD KMTRSHKNVA DDYIHTAACL HSLALEEPTV IKKYLLKVAE LFEKLRKVEG RVSSDEDLKL TELLRYYMLN IEAAKDLLYR RTKALIDYEN SNKALDKARL KSKDVKLAET HQQECCQKFE QLSESAKEEL INFKRKRVAA FRKNLIEMSE LEIKHARNNV SLLQSCIDLF KNN

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Snx5 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 μm filtered
Endotoxin Level:	Protein is endotoxin free.
Grade:	Crystallography grade

Target Details

Target:	SNX5
Alternative Name:	Snx5 (SNX5 Products)

Target Details

Background:

Involved in several stages of intracellular trafficking. Interacts with membranes containing phosphatidylinositol lipids. Acts in part as component of the retromer membrane-deforming SNX-BAR subcomplex. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC). Does not have in vitro vesicle-to-membrane remodeling activity. Involved in retrograde transport of lysosomal enzyme receptor IGF2R. May function as link between endosomal transport vesicles and dynactin. Plays a role in the internalization of EGFR after EGF stimulation. Involved in EGFR endosomal sorting and degradation, the function involves PIP5K1C and is retromer-independent. Together with PIP5K1C facilitates HGS interaction with ubiquitinated EGFR, which initiates EGFR sorting to intraluminal vesicles (ILVs) of the multivesicular body for subsequent lysosomal degradation. Involved in E-cadherin sorting and degradation, inhibits PIP5K1C-mediated E-cadherin degradation (By similarity). Plays a role in macropinocytosis (PubMed:18854019). {ECO:0000250|UniProtKB:Q9Y5X3, ECO:0000269|PubMed:18854019}.

Molecular Weight:

47.6 kDa Including tag.

UniProt:

Q9D8U8

Application Details

An	plication	Notes:
\neg	piication	INOICS.

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.

Comment:

Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.

Handling

Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process