antibodies -online.com





Chemerin Protein (AA 21-156) (His tag)





Go to Product page

()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Overview	
Quantity:	1 mg
Target:	Chemerin (RARRES2)
Protein Characteristics:	AA 21-156
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Chemerin protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA, Crystallization (Crys)
Product Details	
Sequence:	EPELSETQRR SLQVALEEFH KHPPVQLAFQ EIGVDRAEEV LFSAGTFVRL EFKLQQTNCP
	KKDWKKPECT IKPNGRRRKC LACIKMDPKG KILGRIVHCP ILKQGPQDPQ ELQCIKIAQA
	GEDPHGYFLP GQFAFS
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	Made in Germany - from design to production - by highly experienced protein experts.
	Mouse Rarres2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process
	to ensure crystallization grade.
	State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made to order protein and will be made for the first time for your order. Our
	experts in the lab will ensure that you receive a correctly folded protein.

made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Target:

Crystallography grade

Chemerin (RARRES2)

Target Details

ranget.	onement (10 title 2)
Alternative Name:	Rarres2 (RARRES2 Products)
Background:	Adipocyte-secreted protein (adipokine) that regulates adipogenesis, metabolism and
	inflammation through activation of the chemokine-like receptor 1 (CMKLR1). Its other ligands
	include G protein-coupled receptor 1 (GPR1) and chemokine receptor-like 2 (CCRL2). Positively
	regulates adipocyte differentiation, modulates the expression of adipocyte genes involved in

lipid and glucose metabolism and might play a role in angiogenesis, a process essential for the expansion of white adipose tissue. Also acts as a proinflammatory adipokine, causing an increase in secretion of proinflammatory and prodiabetic adipokines, which further impair adipose tissue metabolic function and have negative systemic effects including impaired insulin sensitivity, altered glucose and lipid metabolism, and a decrease in vascular function in other tissues. Can have both pro- and anti-inflammatory properties depending on the modality of enzymatic cleavage by different classes of proteases. Acts as a chemotactic factor for leukocyte populations expressing CMKLR1, particularly immature plasmacytoid dendritic cells, but also immature myeloid DCs, macrophages and natural killer cells. Exerts an anti-inflammatory role by preventing TNF/TNFA-induced VCAM1 expression and monocytes adhesion in vascular endothelial cells. The effect is mediated via inhibiting activation of NF-kappa-B and CRK/p38 through stimulation of AKT1/NOS3 signaling and nitric oxide production. Exhibits an antimicrobial function in the skin. {ECO:0000269|PubMed:17635925, ECO:0000269|PubMed:18242188}.

Molecular Weight:	16.5 kDa Including tag.
UniProt:	Q9DD06
Pathways:	Brown Fat Cell Differentiation

Application Details

· · · · · · · · · · · · · · · · · · ·	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.

Handling

Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process