

Datasheet for ABIN3137290 CDC20 Protein (AA 1-499) (His tag)



[Go to Product page](#)

1 Image

Overview

| | |
|-------------------------------|--|
| Quantity: | 1 mg |
| Target: | CDC20 |
| Protein Characteristics: | AA 1-499 |
| Origin: | Mouse |
| Source: | Insect Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CDC20 protein is labelled with His tag. |
| Application: | Western Blotting (WB), ELISA, SDS-PAGE (SDS), Crystallization (Crys) |

Product Details

Sequence: MAQVFVESDL HSLQLDAPI PNAPVARWQR KAKEATGPAP SPMRAANRSH SAGRTPGRTP
GKSSSKVQTT PSKPGGDRFI PQRSASQMEV ASFLLSKENQ PEDRGTPTKK EHQAWSLNL
NGFDVEEAKI LRLSGKPQNA PEGYQNRLKV LYSQKATPGS SRKTCRYIPS LPDRILDAPE
IRNDYYLNLV DWSSGNVLAV ALDNSVYLWN AGSGDILQLL QMEQPGDYIS SVAWIKEGNY
LAVGTSNAEV QLWDVQQQKR LRNMTSHSAR VSSLSWNSYI LSSGSRSGHI HHHDV RVAEH
HVATLSGHSQ EVCGLRWAPD GRHLASGGND NIVNVWPSGP GESGWAPLQT FTQHQA VKA
VAWCPWQSN I LATGGGTSDR HIRIWNVCSG ACLSAVDVHS QVCSILWSPH YKELISGHGF
AQNQLVIWKY PTMAKVAELK GHTARVLGLT MSPDGATVAS AAADETLRLW RCFEMDPALR
REREKASVAK SSLIHQGIR

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Product Details

- Characteristics:
- Made in Germany - from design to production - by highly experienced protein experts.
 - Mouse Cdc20 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
 - State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

- Purification:
- Two step purification of proteins expressed in baculovirus infected SF9 insect cells:
1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility: 0.22 µm filtered

Endotoxin Level: Protein is endotoxin free.

Grade: Crystallography grade

Target Details

| | |
|-------------------|--|
| Target: | CDC20 |
| Alternative Name: | Cdc20 (CDC20 Products) |
| Background: | Required for full ubiquitin ligase activity of the anaphase promoting complex/cyclosome (APC/C) and may confer substrate specificity upon the complex. Is regulated by MAD2L1: in metaphase the MAD2L1-CDC20-APC/C ternary complex is inactive and in anaphase the CDC20-APC/C binary complex is active in degrading substrates. The CDC20-APC/C complex positively regulates the formation of synaptic vesicle clustering at active zone to the presynaptic membrane in postmitotic neurons. CDC20-APC/C-induced degradation of NEUROD2 induces presynaptic differentiation. {ECO:0000269 PubMed:19900895}. |
| Molecular Weight: | 55.8 kDa Including tag. |
| UniProt: | Q9JJ66 |

Application Details

| | |
|--------------------|---|
| Application Notes: | In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though. |
| Comment: | Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest. |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|--|
| Format: | Liquid |
| Buffer: | 100 mM NaCl, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. |
| Expiry Date: | Unlimited (if stored properly) |



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process