# antibodies .- online.com





# DARC Protein (AA 1-334) (rho-1D4 tag)



## **Image**



Go to Product page

( )	1 /	$\sim$	KI /	110	Νę
	1//	$\vdash$	I \/	1 ←	٠// ٢

Quantity:	1 mg
Target:	DARC
Protein Characteristics:	AA 1-334
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This DARC protein is labelled with rho-1D4 tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

### **Product Details**

## Sequence:

MGNCLYPVET LSLDKNGTQF TFDSWNYSFE DNYSYELSSD YSLTPAAPCY SCNLLDRSSL
PFFMLTSVLG MLASGSILFA ILRPFFHWQI CPSWPILAEL AVGSALFSIA VPILAPGLHS
AHSTALCNLG YWVWYTSAFA QALLIGCYAC LNPRLNIGQL RGFTLGLSVG LWGAAALSGL
PVALASDVYN GFCTFPSSRD MEALKYTHYA ICFTIFTVLP LTLLAAKGLK IALSKGPGPW
VSVLWIWFIF WWPHGMVLIF DALVRSKTVL LYTCQSQKIL DAMLNVTEAL SMLHCVATPL

LLALFCHQTT RRSLSSLSLP TRQASQMDAL AGKS

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

#### Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Mouse Ackr1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

#### Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

- 1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
- 2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.	
Sterility:	0.22 µm filtered	
Endotoxin Level:	Protein is endotoxin-free.	
Grade:	Crystallography grade	

## **Target Details**

Target: DARC


# **Target Details**

Alternative Name:	Ackr1 (DARC Products)		
Background:	Atypical chemokine receptor that controls chemokine levels and localization via high-affinity		
	chemokine binding that is uncoupled from classic ligand-driven signal transduction cascades,		
	resulting instead in chemokine sequestration, degradation, or transcytosis. Also known as		
	interceptor (internalizing receptor) or chemokine-scavenging receptor or chemokine decoy		
	receptor. Has a promiscuous chemokine-binding profile, interacting with inflammatory		
	chemokines of both the CXC and the CC subfamilies but not with homeostatic chemokines.		
	Acts as a receptor for chemokines including CCL2, CCL5, CCL7, CCL11, CCL13, CCL14, CCL17,		
	CXCL5, CXCL6, IL8/CXCL8, CXCL11, GRO, RANTES, MCP-1 and TARC. May regulate chemokine		
	bioavailability and, consequently, leukocyte recruitment through two distinct mechanisms:		
	when expressed in endothelial cells, it sustains the abluminal to luminal transcytosis of tissue-		
	derived chemokines and their subsequent presentation to circulating leukocytes, when		
	expressed in erythrocytes, serves as blood reservoir of cognate chemokines but also as a		
	chemokine sink, buffering potential surges in plasma chemokine levels (By similarity).		
	{ECO:0000250}.		
Molecular Weight:	37.9 kDa Including tag.		
UniProt:	Q9QUI6		
Application Details			
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies		
	as well. As the protein has not been tested for functional studies yet we cannot offer a gurantee		
	though.		
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the		
	recombinant protein with the default tag will be insoluble our protein lab may suggest a higher		
	molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible		
	options with you in detail to assure that you receive your protein of interest.		
Restrictions:	For Research Use only		
Handling			
Handling Format:	Liquid		
	Liquid  100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.		

# Handling

Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

# **Images**



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process