

Datasheet for ABIN3137421

**AMFR Protein (AA 1-643) (rho-1D4 tag)**[Go to Product page](#)**1** Image

## Overview

Quantity:	1 mg
Target:	AMFR
Protein Characteristics:	AA 1-643
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This AMFR protein is labelled with rho-1D4 tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

## Product Details

Sequence:	MPLLFLERFP WPSLRITYTGL SGLALLGTIV SAYRALSQPE DSGGEPEPLT APLQPEALAP ARLTAGGPRA RDVAQYLLSD SLFVWVLVNT ACCVLMLVAK LIQCIVFGPL RVSERQHLKD KFWNFIFYKF IFIFGVLNVQ TVEEVVMWCL WFAGLVFLHL MVQLCKDRFE YLSFSPTTPM SSHGRVLSLL IAMLLSCCGL AVVCCVTGYT HGMHTLAFMA AESLLVTVRT AHVILRYVIH LWDLNHEGTW EGKGTIVVYTT DFMELALLS LDLMHHIHL LFGNIWLSMA SLVIFMQLRY LFHEVQRRIR RHKNYLRVVG NMEARFAVAT PEELAVNND CAICWDSMQA ARKLPCGHLF HNSCLRSWLE QDTSCPTCRM SLNIADGSRA REDHQGENLD ENLVPVAAAE GRPRLNQHNH FFHFDGSRIA SWLPFSVEV MHTTNILGIT QASNSQLNAM AHQIQEMFPQ VPYHLVLQDL QMTRSVEITT DNILEGRIQV PFPTQRSDSL RPALNSPVER PSPDLEESEA SVQTERVPLD LSPRLEETLD FSEVELEPIE VEDFEARGSR FSKSADERQR MLVQRKDDL QQARKRFLNK SSEDDGASER LLPSEGTSST PVTLLRRMLA AAAERRLQRQ RTT
-----------	--

**Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a**

### **special request, please contact us.**

#### Characteristics:

- Made in Germany - from design to production - by highly experienced protein experts.
- Mouse Amfr Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

#### Purification:

Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

1. Membrane proteins are fractionated by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
3. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

#### Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

#### Sterility:

0.22 µm filtered

#### Endotoxin Level:

Protein is endotoxin-free.

## Product Details

Grade: Crystallography grade

## Target Details

Target: AMFR

Alternative Name: Amfr ([AMFR Products](#))

Background: E3 ubiquitin-protein ligase that mediates the polyubiquitination of a number of proteins such as CD3D, CYP3A4, CFTR and APOB for proteasomal degradation. Component of a VCP/p97-AMFR/gp78 complex that participates in the final step of endoplasmic reticulum-associated degradation (ERAD). The VCP/p97-AMFR/gp78 complex is involved in the sterol-accelerated ERAD degradation of HMGCR through binding to the HMGCR-INSIG complex at the ER membrane and initiating ubiquitination of HMGCR. The ubiquitinated HMGCR is then released from the ER by the complex into the cytosol for subsequent destruction. Also acts as a scaffold protein to assemble a complex that couples ubiquitination, retranslocation and deglycosylation. Mediates tumor invasion and metastasis (By similarity). {ECO:0000250, ECO:0000269|PubMed:16987818, ECO:0000269|PubMed:18216283}.

Molecular Weight: 74.3 kDa Including tag.

UniProt: [Q9R049](#)

Pathways: [ER-Nucleus Signaling](#)

## Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

## Handling

Handling Advice:	Avoid repeated freeze-thaw cycles.
------------------	------------------------------------

Storage:	-80 °C
----------	--------

Storage Comment:	Store at -80°C.
------------------	-----------------

Expiry Date:	Unlimited (if stored properly)
--------------	--------------------------------

## Images



**Image 1.** „Crystallography Grade“ protein due to multi-step, protein-specific purification process