antibodies .- online.com





JNK2 Protein (AA 1-423) (Strep Tag)



Overview

Quantity:	1 mg
Target:	JNK2 (MAPK9)
Protein Characteristics:	AA 1-423
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This JNK2 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Sequence: MSDSKSDGQF YSVQVADSTF TVLKRYQQLK PIGSGAQGIV CAAFDTVLGI NVAVKKLSRP

FQNQTHAKRA YRELVLLKCV NHKNIISLLN VFTPQKTLEE FQDVYLVMEL MDANLCQVIH

MELDHERMSY LLYQMLCGIK HLHSAGIIHR DLKPSNIVVK SDCTLKILDF GLARTACTNF

MMTPYVVTRY YRAPEVILGM GYKENVDIWS VGCIMAEMVL HKVLFPGRDY IDQWNKVIEQ

LGTPSAEFMK KLQPTVRNYV ENRPKYPGIK FEELFPDWIF PSESERDKIK TSQARDLLSK

MLVIDPDKRI SVDEALRHPY ITVWYDPAEA EAPPPQIYDA QLEEREHAIE EWKELIYKEV

MDWEERSKNG VKDQPSDAAV SSKATPSQSS SINDISSMST EHTLASDTDS SLDASTGPLE GCR

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression

system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics: Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Product Details	
Purity:	≥ 80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Target Details	
Target:	JNK2 (MAPK9)
Alternative Name:	Mapk9 (MAPK9 Products)
Background:	Mitogen-activated protein kinase 9 (MAP kinase 9) (MAPK 9) (EC 2.7.11.24) (Stress-activated
	protein kinase JNK2) (c-Jun N-terminal kinase 2),FUNCTION: Serine/threonine-protein kinase
	involved in various processes such as cell proliferation, differentiation, migration,
	transformation and programmed cell death. Extracellular stimuli such as pro-inflammatory
	cytokines or physical stress stimulate the stress-activated protein kinase/c-Jun N-terminal
	kinase (SAP/JNK) signaling pathway. In this cascade, two dual specificity kinases
	MAP2K4/MKK4 and MAP2K7/MKK7 phosphorylate and activate MAPK9/JNK2. In turn,
	MAPK9/JNK2 phosphorylates a number of transcription factors, primarily components of AP-1
	such as JUN and ATF2 and thus regulates AP-1 transcriptional activity. In response to oxidative
	or ribotoxic stresses, inhibits rRNA synthesis by phosphorylating and inactivating the RNA
	polymerase 1-specific transcription initiation factor RRN3. Promotes stressed cell apoptosis by
	phosphorylating key regulatory factors including TP53 and YAP1. In T-cells, MAPK8 and MAPK9
	are required for polarized differentiation of T-helper cells into Th1 cells. Upon T-cell receptor
	(TCR) stimulation, is activated by CARMA1, BCL10, MAP2K7 and MAP3K7/TAK1 to regulate
	JUN protein levels. Plays an important role in the osmotic stress-induced epithelial tight-
	junctions disruption. When activated, promotes beta-catenin/CTNNB1 degradation and inhibits
	the canonical Wnt signaling pathway. Participates also in neurite growth in spiral ganglion

neurons. Phosphorylates the CLOCK-BMAL1 heterodimer and plays a role in the regulation of the circadian clock (PubMed:22441692). Phosphorylates POU5F1, which results in the inhibition of POU5F1's transcriptional activity and enhances its proteasomal degradation

 $(PubMed:29153991).\ \{ECO:0000269|PubMed:10811224,\ ECO:0000269|PubMed:11562351,\ PubMed:29153991\}.\ \{ECO:0000269|PubMed:10811224,\ ECO:0000269|PubMed:11562351,\ PubMed:29153991\}.$

ECO:0000269|PubMed:16973441, ECO:0000269|PubMed:21554942, ECO:0000269|PubMed:22441692, ECO:0000269|PubMed:29153991,

200.0000207| 4511164.22 111052, 200.0000207| 4511164.25100551

ECO:0000269|PubMed:9806643}.

Molecular Weight: 48.2 kDa

UniProt: Q9WTU6

Pathways: MAPK Signaling, WNT Signaling, TLR Signaling, Fc-epsilon Receptor Signaling Pathway,

Activation of Innate immune Response, Cellular Response to Molecule of Bacterial Origin,
Positive Regulation of Endopeptidase Activity, Hepatitis C, Toll-Like Receptors Cascades, BCR
Signaling, S100 Proteins

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's
Restrictions:	needed is the DNA that codes for the desired protein! For Research Use only
Handling	To research osciony
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)