



[Go to Product page](#)

Datasheet for ABIN3137587

SLC7A5 Protein (AA 1-512) (Strep Tag)

Overview

Quantity:	1 mg
Target:	SLC7A5
Protein Characteristics:	AA 1-512
Origin:	Mouse
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLC7A5 protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA

Product Details

Sequence: MAVAGAKRRA VATPAAAAE EERQAREKML EARRGDGADP EGEVTLQRN ITLLNGVAII
VGTIIGSGIF VTPTGVLKEA GSPGLSLVW AVCGVFSIVG ALCYaelGTT ISKSGGDYAY
MLEVYGS LPA FLKLWIELLI IRPSSQYIVA LVFATYLLKP VFPTCPVPEE AAKLVACL CV
LLLTAVNCYS VKAATRVQDA FAAAKLLALA LIILLGFIQM GKDMGQGDAS NLQKLSFEG
TNLDVGNIVL ALYSGLFAYG GWNYLNFVTE EMINPYRNLP LAIIISLPV TLVYVLTNLA
YFTTLSTNQMLTSEAVAVDF GNYHLGVMSW IIPV FVGLSC FGSVNGSLFT SSRLFFVGSR
EGHLPSVLSM IHPQLLTPVP SLVFTCIMTL MYAFSRDIFS IINFFSFFNW LCVALAIIGM
MWLRFKKPEL ERPIKVN LAL PVFFILACL F LIAVSFWKTP MECGIGFAI LSGLPVYFFG
VWWKNKPKWI LQAIFSVTVL CQKLMQVVPQ ET

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and

Product Details

Western blot.

Purity: $\geq 80\%$ as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Endotoxin Level: Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

Target Details

Target: SLC7A5

Alternative Name: Slc7a5 ([SLC7A5 Products](#))

Background: Large neutral amino acids transporter small subunit 1 (4F2 light chain) (4F2 LC) (4F2LC) (L-type amino acid transporter 1) (LAT1) (Solute carrier family 7 member 5),FUNCTION: The heterodimer with SLC3A2 functions as a sodium-independent, high-affinity transporter that mediates uptake of large neutral amino acids such as phenylalanine, tyrosine, histidine, methionine, tryptophan, valine, isoleucine and alanine (By similarity). The heterodimer with SLC3A2 mediates the uptake of L-DOPA and leucine (PubMed:9915839, PubMed:11011012). Functions as an amino acid exchanger (By similarity). May play a role in the transport of L-DOPA across the blood-brain barrier (PubMed:11011012). May act as the major transporter of tyrosine in fibroblasts (By similarity). May mediate blood-to-retina L-leucine transport across the inner blood-retinal barrier (By similarity). Can mediate the transport of thyroid hormones diiodothyronine (T2), triiodothyronine (T3) and thyroxine (T4) across the cell membrane. When associated with LAPT4B, the heterodimer formed by SLC3A2 and SLC7A5 is recruited to lysosomes to promote leucine uptake into these organelles, and thereby mediates mTORC1 activation. Involved in the uptake of toxic methylmercury (MeHg) when administered as the L-cysteine or D,L-homocysteine complexes. Involved in the cellular activity of small molecular weight nitrosothiols, via the stereoselective transport of L-nitrosocysteine (L-CNSO) across the membrane (By similarity). {ECO:0000250|UniProtKB:Q01650, ECO:0000250|UniProtKB:Q63016, ECO:0000269|PubMed:11011012, ECO:0000269|PubMed:9915839}.

Molecular Weight: 55.9 kDa

UniProt: [Q9Z127](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Application Details

Comment:	<p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)