

Datasheet for ABIN3137587

SLC7A5 Protein (AA 1-512) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	SLC7A5
Protein Characteristics:	AA 1-512
Origin:	Mouse
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLC7A5 protein is labelled with Strep Tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA

Product Details

Brand:	AliCE®
Sequence:	<p>MAVAGAKRRA VATPAAAAAE EERQAREKML EARRGDGADP EGEGVTLQRN ITLLNGVAII VGTIIGSGIF VTPTGVLKEA GSPGLSLVW AVCGVFSIVG ALCYaelGTT ISKSGGDYAY MLEVYGSLPA FLKLWIELLI IRPSSQYIVA LVFATYLLKP VFPTCPVPEE AAKLVACLCV LLLTAVNCYS VKAATRVQDA FAAAKLLALA LIILLGFIQM GKDMGQGDAS NLQQKLSFEG TNLDVGNIVL ALYSGLFAYG GWNYLNFVTE EMINPYRNLP LAIIISLPV TLVYVLTNLA YFTTLSTNQM LTSEAVAVDF GNYHLGVMSW IIPVFVGLSC FGSVNGSLFT SSRLFFVGSR EGHLPSVL SM IHPQLLTPVP SLVFTCIMTL MYAFSRDIFS IINFFSFFNW LCVALAIIGM MWLRFKKPEL ERPIKVN LAL PVFFILACLF LIAVSFWKTP MECGIGFAI LSGLPVYFFG VWWKNKPKWI LQAIFSVTVL CQKLMQVVPQ ET</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you</p>

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade:

custom-made

Target Details

Target:	SLC7A5
Alternative Name:	Slc7a5 (SLC7A5 Products)
Background:	<p>Large neutral amino acids transporter small subunit 1 (4F2 light chain) (4F2 LC) (4F2LC) (L-type amino acid transporter 1) (LAT1) (Solute carrier family 7 member 5),FUNCTION: The heterodimer with SLC3A2 functions as a sodium-independent, high-affinity transporter that mediates uptake of large neutral amino acids such as phenylalanine, tyrosine, histidine, methionine, tryptophan, valine, isoleucine and alanine (By similarity). The heterodimer with SLC3A2 mediates the uptake of L-DOPA and leucine (PubMed:9915839, PubMed:11011012). Functions as an amino acid exchanger (By similarity). May play a role in the transport of L-DOPA across the blood-brain barrier (PubMed:11011012). May act as the major transporter of tyrosine in fibroblasts (By similarity). May mediate blood-to-retina L-leucine transport across the inner blood-retinal barrier (By similarity). Can mediate the transport of thyroid hormones diiodothyronine (T2), triiodothyronine (T3) and thyroxine (T4) across the cell membrane. When associated with LAPT4B, the heterodimer formed by SLC3A2 and SLC7A5 is recruited to lysosomes to promote leucine uptake into these organelles, and thereby mediates mTORC1 activation. Involved in the uptake of toxic methylmercury (MeHg) when administered as the L-cysteine or D,L-homocysteine complexes. Involved in the cellular activity of small molecular weight nitrosothiols, via the stereoselective transport of L-nitrosocysteine (L-CNSO) across the membrane (By similarity). {ECO:0000250 UniProtKB:Q01650, ECO:0000250 UniProtKB:Q63016, ECO:0000269 PubMed:11011012, ECO:0000269 PubMed:9915839}.</p>
Molecular Weight:	55.9 kDa
UniProt:	Q9Z127

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	<p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the</p>

Application Details

mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.
Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months