

Datasheet for ABIN3137615
DEAF1 Protein (AA 1-566) (His tag)



[Go to Product page](#)

1 Image

Overview

Quantity:	1 mg
Target:	DEAF1
Protein Characteristics:	AA 1-566
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This DEAF1 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

Product Details

Sequence: MEDSDSAAKQ LGLAEAAVA AAAAVAAAAA AAAESEAEPP VLSRDEDSEE DADSEAERET
RRVTAVAVMA AESGHMDMGT EALPSPDEAA AAAAFAEVT TVTVANVGSS ADNVTTSVA
NAASISGHVL SGRTALQIGD SLNTEKATLI VVHTDGSIVE TTGLKGPAAP LTPGPQSPPT
PLAPGQEKGG TKYNWDPVSY DSELPVRCRN ISGTLYKSRL GSGGRGRCIK QGENWYSPT
FEAMAGRASS KDWKRSIRYA GRPLQCLIQD GILNPHAASC TCAACDDMT LSGPVRLFVP
YKRRKKENEL PTTVPKKDSP KNITLLPATA ATTFTVTPSG QITTSALTF DRASTVEATA
VISESPAQGD VFAGATVQEA GVQPPCRVGH PEPHYPGYQD SCQIAPFPEA ALPTSHPKIV
LTSLPALAVP PSTPTKAVSP TVVSGLEMSE HRSWLYLEEM VNSLLNTAQQ LKTLFEQAKQ
ASSCREAAVT QARMQVDTER KEQSCVNCGR EAMSECTGCH KVNVCSTFCQ RKDWKDHQHV
CGQSASVTVQ ADDVHVEESV IEKVAV

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Product Details

- Characteristics:
- Made in Germany - from design to production - by highly experienced protein experts.
 - Mouse Deaf1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
 - State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

-
- Purification:
- Two step purification of proteins expressed in baculovirus infected SF9 insect cells:
1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility: 0.22 µm filtered

Endotoxin Level: Protein is endotoxin free.

Grade: Crystallography grade

Target Details

Target:	DEAF1
Alternative Name:	Deaf1 (DEAF1 Products)
Background:	<p>Transcription factor that binds to sequence with multiple copies of 5'-TTC[CG]G-3' present in its own promoter and that of the HNRPA2B1 gene. Down-regulates transcription of these genes. Binds to the retinoic acid response element (RARE) 5'-AGGGTTCACCGAAAGTTCA-3'. Activates the proenkephalin gene independently of promoter binding, probably through protein-protein interaction (By similarity). Regulates epithelial cell proliferation and side-branching in the mammary gland. Required for neural tube closure and skeletal patterning. Controls the expression of peripheral tissue antigens in pancreatic lymph nodes. Isoform 1 displays greater transcriptional activity than isoform 2. Isoform 2 may inhibit transcriptional activity of isoform 1 by interacting with it and retaining it in the cytoplasm. Transcriptional activator of EIF4G3 (By similarity). May also involved in behavior (PubMed:24726472).</p> <p>{ECO:0000250 UniProtKB:O75398, ECO:0000269 PubMed:14966286, ECO:0000269 PubMed:18826651, ECO:0000269 PubMed:19668219, ECO:0000269 PubMed:24726472}.</p>
Molecular Weight:	60.6 kDa Including tag.
UniProt:	Q9Z1T5
Pathways:	Tube Formation

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCl, 20 mM HEPES, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: Unlimited (if stored properly)

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process