

Datasheet for ABIN3137621

CHEK2 Protein (AA 1-546) (Strep Tag)



Overview

Quantity:	250 μg	
Target:	CHEK2	
Protein Characteristics:	AA 1-546	
Origin:	Mouse	
Source:	Cell-free protein synthesis (CFPS)	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This CHEK2 protein is labelled with Strep Tag.	
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA	

Product Details	
Brand:	AliCE®
Sequence:	MKSHHQSHSS TSSKAHDSAS CSQSQGGFSQ PQGTPSQLHE LSQYQGSSSS STGTVPSSSQ
	SSHSSSGTLS SLETVSTQEL CSIPEDQEPE EPGPAPWARL WALQDGFSNL DCVNDNYWFG
	RDKSCEYCFD GPLLRRTDKY RTYSKKHFRI FREMGPKNCY IVYIEDHSGN GTFVNTELIG
	KGKRCPLSNN SEIALSLCRN KVFVFFDLTV DDQSVYPKEL RDEYIMSKTL GSGACGEVKM
	AFERKTCQKV AIKIISKRRF ALGSSREADT APSVETEIEI LKKLNHPCII KIKDVFDAED YYIVLELMEG
	GELFDRVVGN KRLKEATCKL YFYQMLVAVQ YLHENGIIHR DLKPENVLLS SQEEDCLIKI
	TDFGQSKILG ETSLMRTLCG TPTYLAPEVL VSNGTAGYSR AVDCWSLGVI LFICLSGYPP
	FSEHKTQVSL KDQITSGKYN FIPEVWTDVS EEALDLVKKL LVVDPKARLT TEEALNHPWL
	QDEYMKKKFQ DLLVQEKNSV TLPVAPAQTS SQKRPLELEV EGMPSTKRLS VCGAVL
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- · Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	CHEK2
Alternative Name:	Chek2 (CHEK2 Products)
Background:	Serine/threonine-protein kinase Chk2 (EC 2.7.11.1) (CHK2 checkpoint homolog) (Checkpoint
	kinase 2),FUNCTION: Serine/threonine-protein kinase which is required for checkpoint-mediated
	cell cycle arrest, activation of DNA repair and apoptosis in response to the presence of DNA
	double-strand breaks. May also negatively regulate cell cycle progression during unperturbed
	cell cycles. Following activation, phosphorylates numerous effectors preferentially at the
	consensus sequence [L-X-R-X-X-S/T]. Regulates cell cycle checkpoint arrest through
	phosphorylation of CDC25A, CDC25B and CDC25C, inhibiting their activity. Inhibition of CDC25
	phosphatase activity leads to increased inhibitory tyrosine phosphorylation of CDK-cyclin
	complexes and blocks cell cycle progression. May also phosphorylate NEK6 which is involved
	in G2/M cell cycle arrest. Regulates DNA repair through phosphorylation of BRCA2, enhancing
	the association of RAD51 with chromatin which promotes DNA repair by homologous
	recombination. Also stimulates the transcription of genes involved in DNA repair (including
	BRCA2) through the phosphorylation and activation of the transcription factor FOXM1.
	Regulates apoptosis through the phosphorylation of p53/TP53, MDM4 and PML.
	Phosphorylation of p53/TP53 at 'Ser-20' by CHEK2 may alleviate inhibition by MDM2, leading to
	accumulation of active p53/TP53. Phosphorylation of MDM4 may also reduce degradation of
	p53/TP53. Also controls the transcription of pro-apoptotic genes through phosphorylation of
	the transcription factor E2F1. Tumor suppressor, it may also have a DNA damage-independent
	function in mitotic spindle assembly by phosphorylating BRCA1. Its absence may be a cause of
	the chromosomal instability observed in some cancer cells. Promotes the CCAR2-SIRT1
	association and is required for CCAR2-mediated SIRT1 inhibition (By similarity).
	{ECO:0000250 UniProtKB:096017, ECO:0000269 PubMed:12192050,
	ECO:0000269 PubMed:25619829}.
Molecular Weight:	61.1 kDa
UniProt:	Q9Z265
Pathways:	p53 Signaling, Apoptosis, Cell Division Cycle
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.

Application Details

_			
\cap	m	m	nt:

ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months