ANTIBODIES ONLINE

Datasheet for ABIN3197487 anti-Isocitrate Dehydrogenase antibody

1 Image

3 Publications



Overview

Quantity:	50 µg
Target:	Isocitrate Dehydrogenase (IDH)
Reactivity:	Arabidopsis thaliana, Pisum sativum, Potato, Tomato, Zea mays
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB)
Product Details	
Immunogen:	KLH-conjugated peptide 1 and peptide 2 conserved in all higher plants mitochondrial, NAD
Immunogen:	KLH-conjugated peptide 1 and peptide 2 conserved in all higher plants mitochondrial, NAD dependent isocitrate dehydrogenase subunits including Arabidopsis thaliana IDH-I Q8LFC0,
Immunogen:	KLH-conjugated peptide 1 and peptide 2 conserved in all higher plants mitochondrial, NAD dependent isocitrate dehydrogenase subunits including Arabidopsis thaliana IDH-I Q8LFC0, At4g35260 and IDH-II P93032, At4g17130
Immunogen: Cross-Reactivity (Details):	KLH-conjugated peptide 1 and peptide 2 conserved in all higher plants mitochondrial, NAD dependent isocitrate dehydrogenase subunits including Arabidopsis thaliana IDH-I Q8LFC0, At4g35260 and IDH-II P93032, At4g17130 No cross-reactivity with: Chlamydomonas reinhardtii
Immunogen: Cross-Reactivity (Details): Predicted Reactivity:	KLH-conjugated peptide 1 and peptide 2 conserved in all higher plants mitochondrial, NAD dependent isocitrate dehydrogenase subunits including Arabidopsis thaliana IDH-I Q8LFC0, At4g35260 and IDH-II P93032, At4g17130 No cross-reactivity with: Chlamydomonas reinhardtii dicots including Brassica napus, Vitis vinifera, monocots including Oryza sativa, Zea mays
Immunogen: Cross-Reactivity (Details): Predicted Reactivity: Characteristics:	KLH-conjugated peptide 1 and peptide 2 conserved in all higher plants mitochondrial, NAD dependent isocitrate dehydrogenase subunits including Arabidopsis thaliana IDH-I Q8LFC0, At4g35260 and IDH-II P93032, At4g17130 No cross-reactivity with: Chlamydomonas reinhardtii dicots including Brassica napus, Vitis vinifera, monocots including Oryza sativa, Zea mays Expected / apparent Molecular Weight of the Antigene: 39 / 45 kDa (Arabidopsis thaliana)
Immunogen: Cross-Reactivity (Details): Predicted Reactivity: Characteristics: Purification:	KLH-conjugated peptide 1 and peptide 2 conserved in all higher plants mitochondrial, NAD dependent isocitrate dehydrogenase subunits including Arabidopsis thaliana IDH-I Q8LFC0, At4g35260 and IDH-II P93032, At4g17130 No cross-reactivity with: Chlamydomonas reinhardtii dicots including Brassica napus, Vitis vinifera, monocots including Oryza sativa, Zea mays Expected / apparent Molecular Weight of the Antigene: 39 / 45 kDa (Arabidopsis thaliana) serum

Target Details

Target:	Isocitrate Dehydrogenase (IDH)
Abstract:	IDH Products
Background:	AGI Code: At4g35260
	Plant NADH dependent isocitrate dehydrogenase enzyme is located in mitochondrial matrix.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN3197487 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Molecular Weight:	This enzyme is classified as an oxidoreductase and its function is to catalyze a reaction in the citric acid cycle, specifically the sequential dehydrogenation and decarboxylation of isocitrate to form a-ketoglutarate. It removes hydrogens from its substrate, isocitrate. In addition to this process, it functions as a decarboxylase, removing a CO2 from the six-carbon substrate to form a five-carbon product mentioned above as a-ketoglutarate. There are two forms of this enzyme NADP+ and NAD+ dependent. expected: 39 kDa, apparent: 45 kDa (Arabidopsis thaliana)
UniProt:	P93032
Application Details	
Application Notes:	Recommended Dilution 1 : 5 000 with standard ECL (WB). Additional Information: Peptide used to elicit this antibody is not conserved in NADPH dependentanzymes, partially conserved across eukaryotic Idh subunits. Some conservationacross bacterial which contain the NAD- dependent form of Idh (as opposed to theNADP-dependent form).
Comment:	cellular [compartment marker] of mitochondrial matrix
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	PBS pH 7.4
Handling Advice:	Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes. Once reconstituted make aliquots to avoid repreated freeze-thaw cycles.
Storage:	-20 °C
Publications	
Product cited in:	Rurek, Woyda-Ploszczyca, Jarmuszkiewicz et al.: "Biogenesis of mitochondria in cauliflower (Brassica oleracea var. botrytis) curds subjected to temperature stress and recovery involves regulation of the complexome, respiratory chain activity," in: Biochimica et biophysica acta , Vol. 1847, Issue 4-5, pp. 399-417, (2015) (PubMed).
	Lee, Lee, Yoo, Duncan, Oh, Lee, Lee, Whelan, Hwang: "Mitochondrial targeting of the Arabidopsis

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN3197487 | 07/25/2024 | Copyright antibodies-online. All rights reserved. F1-ATPase γ-subunit via multiple compensatory and synergistic presequence motifs." in: **The Plant cell**, Vol. 24, Issue 12, pp. 5037-57, (2013) (PubMed).

Szal, Jastrz?bska, Kulka, Lea: "Influence of mitochondrial genome rearrangement on cucumber leaf carbon and nitrogen metabolism." in: **Planta**, Vol. 232, Issue 6, pp. 1371-82, (2010) (PubMed).

Images



Image 1. Figure description: 1: Total extract A. thaliana (20 g protein) 2: Fraction enriched with mitochondria A. thaliana 3: Pure mitochondria A. thaliana 4: Pure mitochondria P. satisum 5: Pure mitochondria S. tuberosum Description of SDS-PAGE experimental conditions: After gel electrophoresis samples have been transferred to nitrocellulose membrane. Blocking has been done in 5% milk powder in TBS followed by incubation with primary antibodies for 1 hour and 30 minutes in RT After incubation with secondary antibodies reaction has been developed using ECL reagent (GE Healthcare) * Band detected at ca. 90 kDa is suspected to be a dimmer of Idh, since this band is depleted upon peptide competition experiment.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN3197487 | 07/25/2024 | Copyright antibodies-online. All rights reserved.