antibodies - online.com







anti-SNX19 antibody (Internal Region)



Image



Overview

Quantity:	100 μg
Target:	SNX19
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This SNX19 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	SNX19
Immunogen:	Peptide with sequence C-TTSASDTPGNSKR, from the internal region of the protein sequence according to NP_055573.2.
Sequence:	TTSASDTPGN SKR
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details

rarget Details	
Target:	SNX19
Alternative Name:	SNX19 (SNX19 Products)
Background:	SNX19, sorting nexin 19, CHET8, DKFZp667I205, KIAA0254
Gene ID:	399979
NCBI Accession:	NP_055573
Application Details	
Application Notes:	Western Blot: Approx.110 kDa band observed in Human Skeletal Muscle lysates (calculated
	MW of 109 kDa according to NP_055573.2). Recommended concentration: 0.3-1 μ g/mL.
	Peptide ELISA: antibody detection limit dilution 1:16000.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum
	albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated
	at 4°C for a few weeks and still remain viable.

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

Image 1. EB08885 (0.3μg/ml) staining of Human Skeletal Muscle lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.