

Datasheet for ABIN336452

Donkey anti-Goat IgG Antibody (TRITC)[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	IgG
Reactivity:	Goat
Host:	Donkey
Clonality:	Polyclonal
Conjugate:	TRITC
Application:	Immunohistochemistry (IHC), Flow Cytometry (FACS)

Product Details

Immunogen:	Donkey antiserum was obtained from animals of US origin and under the care of a registered veterinarian. Country of Origin: Donkey antiserum was obtained from animals of US origin and under the care of a registered veterinarian.
Characteristics:	FLUOROPHORE: Tetramethylrhodamine-5-isothiocyanate (TRITC) Amax = 550 nm, Emax = 570 nm. TRITC/Protein Ratio: Moles TRITC per Mole Antibody
Purification:	Antibody was affinity purified using solid phase Goat IgG

Target Details

Target:	IgG
Abstract:	IgG Products
Target Type:	Antibody

Application Details

Application Notes:	This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. Preservative: 0.05 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Product is photosensitive and should be protected from light.
Storage:	4 °C/-20 °C