

Datasheet for ABIN349633
anti-Interleukin 17a antibody



[Go to Product page](#)

3 Images

1 Publication

Overview

Quantity:	100 µg
Target:	Interleukin 17a (IL17A)
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Interleukin 17a antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Neutralization (Neut), Radioimmunoassay (RIA)

Product Details

Immunogen:	This purified antibody was prepared from whole rabbit serum produced by repeated immunizations with full length recombinant rat IL17-A protein. Immunogentype:Recombinant
Isotype:	IgG
Cross-Reactivity:	Mouse (Murine)
Characteristics:	Concentration Definition: by UV absorbance at 280 nm

Target Details

Target:	Interleukin 17a (IL17A)
Alternative Name:	IL-17A (IL17A Products)
Background:	Rat IL17-A (also known as Interleukin-17, Cytotoxic T-lymphocyte-associated antigen 8 and

Target Details

CTLA-8) is a proinflammatory cytokine member of a six-species family of proteins (IL-17A-17F). Rat IL-17A protein is a homodimer consisting of two 134 amino acids peptides. IL-17A is secreted mainly by activated CD4+ and CD8+ T lymphocytes and acts through its receptor, IL-17R, to induce the expression of many mediators of inflammation, most strikingly, those that are involved in the proliferation, maturation and chemotaxis of neutrophils. Elevated levels of IL-17A have been associated with several conditions, including rheumatoid arthritis, airway inflammation, allograft rejection, inflammatory bowel disease, psoriasis, cancer and multiple sclerosis. There is 58% identity between the amino acid sequence of human and rat IL-17A. Synonyms: Interleukin-17, Cytotoxic T-lymphocyte-associated antigen 8 and CTLA-8

Gene ID: 301289

UniProt: [Q61453](#)

Application Details

Application Notes: This purified antibody has been tested for use in ELISA and western blotting. Reactivity is also expected in neutralizations, radioimmunoassay and immunohistochemistry. The endotoxin content is estimated to be <10 pg/μl by the LAL method. The recombinant immunogen is a non-glycosylated homodimer joined by disulfide bonds having a molecular mass of 30.0 kDa. By western blot from a reducing gel expect a band approximately 15.0 kDa in size corresponding to rat IL17-A protein in the appropriate cell lysate or extract. Specific conditions for reactivity should be optimized by the end user.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Restore with deionized water (or equivalent)

Concentration: 1.0 mg/mL

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2


Storage: 4 °C

Publications

Product cited in: Fu, Zhao, Dong, Du, Chen, Wu, Cheng, Du, Liao: "Interleukin-17A contributes to the development of post-operative atrial fibrillation by regulating inflammation and fibrosis in rats with sterile


pericarditis." in: **International journal of molecular medicine**, Vol. 36, Issue 1, pp. 83-92, (2016) ([PubMed](#)).

Images




Western Blotting

Image 1. Expression of atrial fibrillation (AF)-related pro-inflammatory cytokines at 4 days after surgery. (A) Fold mRNA expression of interleukin (IL)-6, IL-1 β , transforming growth factor- β 1 (TGF- β 1) and IL-17A. (B) Protein expression of IL-17A detected by western blot analysis. (C) Quantitative analysis of IL-17A protein expression. * $P < 0.05$ and ** $P < 0.01$ vs. Sham. # $P < 0.05$ and ## $P < 0.01$ vs. IgG2a. Sham, sham-operated rats, SP, rats with sterile pericarditis. - figure provided by CiteAb. Source: PMID25955429



Western Blotting

Image 2. Western blot using anti-IL-17A antibody shows detection of mouse recombinant IL-17A protein (arrowhead, lane 1). Approximately 2 μ g of recombinant protein was loaded onto the gel. Primary antibody was used at a 1:1,000 dilution. The membrane was washed and reacted with a 1:20,000 dilution of 649 conjugated Gt-a-Rabbit IgG. Molecular weight estimation was made by comparison to prestained MW markers indicated at the left. Other detection systems will yield similar results.



Western Blotting

Image 3. Western blot using anti-IL-17A antibody shows detection of rat recombinant IL-17A protein (lane 1). Approximately 2 μ g of recombinant protein was loaded onto the gel. Primary antibody was used at a 1:1,000 dilution. The membrane was washed and reacted with a 1:20,000 dilution of 649 conjugated Gt-a-Rabbit IgG. Molecular weight estimation was made by comparison to prestained MW markers indicated at the left. Other detection systems will yield similar results.