antibodies .- online.com





Datasheet for ABIN350943

anti-Syntaxin 7 antibody



Overview

Quantity:	500 μg
Target:	Syntaxin 7 (STX7)
Reactivity:	Human
Host:	Sheep
Clonality:	Polyclonal
Conjugate:	This Syntaxin 7 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	A synthetic peptide as a part of human Syntaxin-7 (STX7, Syn7, Syntaxin 7) conjugated to an
	immunogenic carrier protein was used as the immunogen.
Isotype:	IgG
Specificity:	Specific for Syntaxin-7.
Cross-Reactivity:	Human
Cross-Reactivity (Details):	Other species not yet tested.
Purification:	IgG
Target Details	
Target:	Syntaxin 7 (STX7)
Alternative Name:	Syntaxin 7 (STX7 Products)

Target Details

Expiry Date:

12 months

Target Details	
Background:	FUNCTION: May be involved in protein trafficking from the plasma membrane to the early endosome (EE) as well as in homotypic fusion of endocytic organelles. Mediates the endocytic trafficking from early endosomes to late endosomes and lysosomes. SUBUNIT: Forms a SNARE complex with VTI1B, STX8 and VAMP8 which functions in the homotypic fusion of late endosomes. Interacts with VPS33A. Found in a complex with VAMP8 and VTI1B in the liver. Part of the SNARE core complex containing VAMP8, STX8 and VTI1B. Interacts with VPS11, VPS16 and VPS18. SUBCELLULAR LOCATION: Early endosome membrane, Single-pass type IV membrane protein.,Syntaxin Family,Syntaxin-7, STX7, Syn7
UniProt:	015400
Pathways:	Synaptic Vesicle Exocytosis
Application Details	
Application Notes:	IHC, WB. A concentration of 10-50 μ g,ml is recommended. The optimal concentration should be determined by the end user. Not yet tested in other applications.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Reconstitute in 500 µL of sterile water. Centrifuge to remove any insoluble material.
Handling Advice:	Avoid freeze and thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles.