antibodies - online.com







anti-PAG1 antibody (AA 97-432, AA 235-280)

Images



\sim					
	1//	⊃r	V/I	Φ\	Λ

Quantity:	50 μg	
Target:	PAG1	
Binding Specificity:	AA 97-432, AA 235-280	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This PAG1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS), Immunohistochemistry (IHC)	

Product Details


Brand:	IHC-plus™	
Immunogen:	Recombinant intracellular fragment (aa 97-432) of human Cbp (PAG).	
	Type of Immunogen: Recombinant protein	
Clone:	MEM-255	
Isotype:	lgG2a	
Recognizes an epitope (aa 235-280) of Csk-binding protein (Cbp) located in the cyto domain, also known as protein associated with glycosphingolipid-enriched microdor (PAG)		
No Cross-Reactivity:	es-Reactivity: Mouse (Murine), Cow (Bovine), Rat (Rattus)	

Product Details		
Purification:	Protein A purified	
Target Details		
Target:	PAG1	
Alternative Name:	PAG1 / PAG (PAG1 Products)	
Background:	Name/Gene ID: PAG1	
	Synonyms: PAG1, Csk-binding protein, CBP, PAG	
Gene ID:	55824	
UniProt:	Q9NWQ8	
Pathways:	p53 Signaling, TCR Signaling, EGFR Signaling Pathway, CXCR4-mediated Signaling Events	
Application Details		
Application Notes:	Approved: Flo (2 μg/mL), IHC, IHC-P (10 μg/mL), WB	
	Usage: Immunohistochemistry: This antibody was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody,	
Comment:	Target Species of Antibody: Human	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, 15 mM sodium azide, approx, pH 7.4.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Avoid repeated freezing and thawing.	
Storage:	4 °C,-20 °C	

Storage Comment:


4°C or -20°C, Avoid freeze-thaw cycles.

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Human Tonsil (formalin-fixed, paraffin-embedded) stained with PAG1 antibody ABIN351517 at 10 ug/ml followed by biotinylated anti-mouse IgG secondary antibody ABIN481714, alkaline phosphatase-streptavidin and chromogen.



Immunohistochemistry

Image 2. Anti-PAG1 / PAG antibody IHC of human tonsil. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody concentration 10 ug/ml.