

Datasheet for ABIN356551

anti-Kdm6b antibody (Middle Region)





Go to Product page

_				
()	ve.	rv/	101	Λ

3 7 3 7 7 3 7 7			
Quantity:	0.4 mL		
Target:	Kdm6b		
Binding Specificity:	Middle Region		
Reactivity:	Human, Mouse		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This Kdm6b antibody is un-conjugated		
Application:	Western Blotting (WB), Enzyme Immunoassay (EIA)		
Product Details			
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide		
	selected from the center region of mouse JMJD3.		
Isotype:	Ig Fraction		
Specificity:	This antibody will recognize JMJD3 (Center).		
Purification:	Protein G Chromatography, eluted with high and low pH buffers and neutralized immediately,		
	followed by dialysis against PBS.		
Target Details			
Target:	Kdm6b		
Alternative Name:	JMJD3 / KDM6B (Kdm6b Products)		

Target Details

Background:	Covalent modification of histones plays critical role in regulating chromatin structure and	
	transcription. While most covalent histone modifications are reversible, only recently has it been	
	established that methyl groups are subject to enzymatic removal from histones. A family of	
	novel JmjC domain-containing histone demethylation (JHDM) enzymes have been identified	
	that perform this specific function. Histone demethylation by JHDM proteins requires cofactors	
	Fe(II) and alpha-ketoglutarate. Family members include JHDM1 (demethylating histone 3 at	
	lysine 36), and JHDM2A as well as JMJD2CH3K9 (both of which demethylate histone 3 at	
	lysine 9). Contributions of histone demethylase activity to tumor development, decreases in cell	
	proliferation, and hormone-dependent transcriptional activation have been observed. Synonyms:	
	Histone demethylase JMJD3, JmjC domain-containing protein 3, Jumonji domain-containing	
	protein 3, KIAA0346, Lysine demethylase 6B	
Gene ID:	216850, 10090	
UniProt:	Q5NCY0	
Pathways:	Warburg Effect	
Application Details		
Application Notes:	ELISA: 1/1,000. Western Blot: 1/250-1/500.	
	Other applications not tested.	
	Optimal dilutions are dependent on conditions and should be determined by the user.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.25 mg/mL	
Buffer:	PBS containing 0.09 % (W/V) Sodium Azide as preservative.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Handling Advice:	Avoid repeated freezing and thawing.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.	




Image 1.