

Datasheet for ABIN356732 anti-PRMT8 antibody (C-Term)

1 Image



Go to Product page

Overview	
Quantity:	0.4 mL
Target:	PRMT8
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRMT8 antibody is un-conjugated
Application:	Western Blotting (WB), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide
	selected from the C-terminal region of human PRMT8.
Isotype:	lg Fraction
Specificity:	This antibody is specific to PRMT8 (C-term).
Purification:	Protein A Chromatography, eluted with high and low pH buffers and neutralized immediately,
	followed by dialysis against PBS.
Target Details	
Target:	PRMT8
Alternative Name:	PRMT8 / HRMT1L3 (PRMT8 Products)

Target Details

rarget betails	
Background:	Protein arginine methylation plays a role in signal transduction, RNA processing, transcriptional regulation, and DNA repair. PRMT8, a protein arginine N-methyltransferase most closely related to PRMT1, methylates the guanidino nitrogens of arginyl residues in some proteins. This protein associates with the plasma membrane following myristoylation and exhibits a brain-specific expression pattern, making it unique within this family of enzymes. Synonyms: HRMT1L4, Heterogeneous nuclear ribonucleoprotein methyltransferase-like protein 4, Protein arginine N-methyltransferase 8
Gene ID:	56341, 9606
UniProt:	Q9NR22
Application Details	
Application Notes:	ELISA: 1/1,000. Western Blot: 1/50-1/100. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS containing 0.09 % (W/V) Sodium Azide as preservative.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at-20 °C for longer.




Image 1.