

Datasheet for ABIN357051

anti-TLR5 antibody (N-Term)

2 Images



Overview

OVEIVIEW	
Quantity:	0.4 mL
Target:	TLR5
Binding Specificity:	N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TLR5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of mouse TLR5.
Isotype:	lg Fraction
Specificity:	This antibody detects TLR5 at N-term.
Purification:	Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS
Target Details	
Target:	TLR5
Alternative Name:	TLR5 (TLR5 Products)

Target Details

Background:	TLR5, a Type I membrane protein belonging to the Toll-like receptor family, participates in the		
	innate immune response to microbial agents. It also plays a role in mediating detection of		
	bacterial flagellins. TLR5 acts via MyD88 and TRAF6, leading to NF-kappa-B activation, cytokine		
	secretion and the inflammatory response. This protein binds to TIRAP and MyD88 via their		
	respective TIR domains TLR5 is highly expressed in liver, and is detected in lung and at very low		
	levels in most other tissues. The TLR5 gene lies in a locus that is associated with susceptibility		
	to Salmonella. Inbred strains of mice can be classified into 3 categories according to their resistance to infection with S.typhimurium: susceptible (BALB/c, C57BL/6, C3H/He), intermediate (DBA/2, C75L) and resistant (A, CBA). The strain MOLF/Ei is highly susceptible to the infection, has an unique TLR5 haplotype and a lower expression of TRL5.Synonyms: TIL3,		
		Toll-like receptor 5, Toll/interleukin-1 receptor-like protein 3	
		Molecular Weight:	97626 Da
		Gene ID:	7100, 10090
	UniProt:	060602	
Pathways:	TLR Signaling, Activation of Innate immune Response, Cellular Response to Molecule of		
	Bacterial Origin, Toll-Like Receptors Cascades		
A 1: 1: D 1:1			
Application Details			
Application Notes:	ELISA 1: 1,000. Western blot 1: 100 - 1: 500. Immunohistochemistry 1: 50 - 1: 100.		
	Other applications not tested.		
	Optimal dilutions are dependent on conditions and should be determined by the user.		
Restrictions:	Optimal dilutions are dependent on conditions and should be determined by the user. For Research Use only		
Restrictions: Handling			
Handling	For Research Use only		
Handling Format:	For Research Use only Liquid		
Handling Format: Concentration:	For Research Use only Liquid 0.25 mg/mL		
Handling Format: Concentration: Buffer:	For Research Use only Liquid 0.25 mg/mL PBS with 0.09 % (W/V) sodium azide		
Handling Format: Concentration: Buffer: Preservative:	For Research Use only Liquid 0.25 mg/mL PBS with 0.09 % (W/V) sodium azide Sodium azide		

Handling

Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer.

Images




Image 1.




Image 2.