antibodies -online.com







anti-ZNF395 antibody (Center)

Images



| Overview | |
|----------------------|--|
| Quantity: | 0.4 mL |
| Target: | ZNF395 |
| Binding Specificity: | Center |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ZNF395 antibody is un-conjugated |
| Application: | Western Blotting (WB), Enzyme Immunoassay (EIA) |
| Product Details | |
| Immunogen: | This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide |
| | selected from the Center region of human HDBP2. |
| Isotype: | Ig Fraction |
| Specificity: | This antibody detects HDBP2(HDRF-2) at center. |
| Purification: | Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by |

Target Details

| Target: | ZNF395 |
|-------------------|--------------------------|
| Alternative Name: | ZNF395 (ZNF395 Products) |

dialysis against PBS

Target Details

| rarget Details | |
|---------------------|---|
| Background: | Papillomavirus Regulatory Factor 1/HDBP2 is a novel transcription factor shuttling between nucleus and cytoplasm and binds to the specific GCCGGCG, which is an essential cis-element for Huntington's disease gene expression. Synonyms: HD-regulating factor 2, HDBP2, Huntington disease gene regulatory region-binding protein 2, PBF, Papillomavirus regulatory factor 1, Papillomavirus-binding factor, Zinc finger protein 395 |
| Molecular Weight: | 54808 Da |
| Gene ID: | 55893, 9606 |
| UniProt: | Q9H8N7 |
| Application Details | |
| Application Notes: | ELISA 1: 1,000. Western blot 1: 50 - 1: 100. Immunohistochemistry 1: 50 - 1: 100. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 0.25 mg/mL |
| Buffer: | PBS with 0.09 % (W/V) sodium azide |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice: | Avoid repeated freezing and thawing. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. |




Image 1.




Image 2.